

10/500, 586
Rec'd PCT/PTO 20 JAN 2006

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 July 2003 (31.07.2003)

PCT

(10) International Publication Number
WO 03/062470 A1

(51) International Patent Classification⁷: C12Q 1/68

(21) International Application Number: PCT/KR03/00131

(22) International Filing Date: 21 January 2003 (21.01.2003)

(25) Filing Language: Korean

(26) Publication Language: English

(30) Priority Data:
10-2002-0004297 24 January 2002 (24.01.2002) KR
10-2002-0011648 5 March 2002 (05.03.2002) KR

(71) Applicant (for all designated States except US):
BIOMEDLAB CORPORATION [KR/KR]; Dong-
sung Bldg., 1-49, Dongsung-dong, Jongro-ku, Seoul
110-510 (KR).

(71) Applicant and

(72) Inventor: **KIM, Bum-Joon** [KR/KR]; A-105 Professor's
apt., 1 Ara 1-dong, Jeju-city, Jeju-do 690-121 (KR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KOOK, Yoon-Ho**

[KR/KR]; 2-1406 Yeoksamhansin Apt., Dogok 1-dong,
Gangnam-gu, Seoul 135-271 (KR). **KIM, Jeong-Mi**
[KR/KR]; 7-208 Bulkwangmiseong apt., Bulkwang
1dong, Eunpyeong-gu, Seoul 122-751 (KR).


(74) Agent: **YOU ME PATENT & LAW FIRM**; Teheran
Bldg., 825-33, Yoksam-dong, Kangnam-ku, Seoul 135-080
(KR).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC,
VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,

[Continued on next page]

(54) Title: PRIMERS FOR AMPLIFYING HSP 65 GENE OF MYCOBACTERIAL SPECIES, HSP 65 GENE FRAGMENTS AND
METHOD OF IDENTIFYING MYCOBACTERIAL SPECIES WITH THE SAME



(57) Abstract: The present invention relates to a pair of primers specific to mycobacterial species, a polynucleotide of an hsp 65 gene fragment, and a method for the identification of mycobacterial species by using the same. More specifically, the 604-bp hsp 65 gene fragment can be applied to identification methods of mycobacteria such as the comparative sequence analysis method, the probe hybridization method, and PCR-RFLP, which can resolve the problems of a conventional identification method based on biochemical characteristics, where the genus mycobacterium covers various species and has a low growth rate, and of the problems of 16s rDNA. Thus, according to the identification method of the present invention, the mycobacterial species can be identified simply, economically, and accurately.

WO 03/062470 A1



SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

PRIMERS FOR AMPLIFYING HSP 65 GENE OF MYCOBACTERIAL SPECIES, HSP 65 GENE FRAGMENTS, AND METHOD OF IDENTIFYING MYCOBACTERIAL SPECIES WITH THE SAME

CROSS REFERENCE TO RELATED APPLICATION

5 The present application is based on Korean patent application No. 2002-004297 filed on January 24, 2002, and Korean patent application No. 2002-0011648 filed on March 5, 2002.

10 **BACKGROUND OF THE INVENTION**

(a) Field of the Invention

The present invention relates to a pair of primers specific to Mycobacterial species, more specifically to a pair of primers that can specifically amplify the hsp 65 gene of mycobacteria, a gene fragment of hsp 65, and an identifying method of Mycobacterial species.

(b) Description of the Related Art

The genus *Mycobacterium* covers a wide range of organisms including obligate species causing serious human and animal disease such as tuberculosis, bovine tuberculosis, and leprosy; opportunistic pathogens; and saprophytic species found in the natural environment. At present, it is known that about 72 species of the genus *mycobacterium* have been reported, of which about 25 species are involved in the human diseases.

Tuberculosis is the largest of the Mycobacterial infections. The Mycobacterial species causing tuberculosis include *M. tuberculosis*, *M. bovis*, *M. africanum*, and *M. microti*, which are classified as *M. tuberculosis* complex (TB complex). *M. tuberculosis* is common and important in causing tuberculosis. Tuberculosis infection decreased because of continuous use of antituberculosis drugs until the end of the 1980s, but in line with the rapid increase of AIDS and Mycobacterium tuberculosis with drug resistance, tuberculosis increased in developed countries in the 1990s. In particular, it has been reported that the death rate due to tuberculosis is the highest among infectious diseases in Korea, claiming about three hundred or more lives per year, because of the increase in the number of street people in the International Monetary Fund era in Korea.

Mycobacteria Other than Mycobacterium tuberculosis (MOTT, or nontuberculous mycobacteria, NTM) causes infection in aged people and immuno-compromised patients, and its clinical manifestation is similar to tuberculosis. The occurrence of MOTT is still lower than tuberculosis in Korea, but it is quite common. It is difficult to determine the pathogenicity from isolate that is separated from a clinical sample. In addition, resistance of MOTT to most anti-tuberculosis drugs and its recurrence rate makes it difficult to treat MOTT infection. It has been reported that MOTT also causes disease in patients who are not immuno-compromised, and that 50% of Mycobacterial infection in the United States is tuberculosis and 50% is MOTT infection over the past 10 years. With the spread of HIV (Human

immunodeficiency virus) infection since the 1980s, MOTT has caused systemic disseminated infection of immuno-compromised patients. Thus, MOTT has been closely watched.

Mycobacterial species have different patterns of resistance to antituberculosis drugs from each other, and thus they are treated by different methods with different drugs (Wolinsky E: Mycobacterial diseases other than tuberculosis. *Clin Infect Dis* 15: 1-10, 1992). Accordingly, Mycobacteria need to be differentiated and identified on a species level.

A biochemical method for identifying Mycobacterial species is laborious and time-consuming due to the slow growing rate of Mycobacteria. A cell wall lipid analyzing method using High-performance Lipid Chromatography (HPLC) and Thin Layer Lipid Chromatography (TLC) is difficult to perform and is costly, and thus it is carried out on a small laboratory scale. The use of conventional identifying methods has a disadvantage in that it takes a great deal of time to perform due to the slow growing rate of the Mycobacteria (about 2-3 months for slow-growing mycobacteria). Thus, the treatment of Mycobacterial infection can be delayed (Nolte FS, Metchock B: *Mycobacterium*, In Murray PR, Baron EJ, Pfaller MA, Tenover FC, Tenover FC, Tenover FC, Yolken RH (eds.), Manual of clinical microbiology. American Society for Microbiology, Washington, D.C. 400-437, 1995.).

16s rDNA is commonly used as a chronometer molecule for identification of the Mycobacterial species with a molecular biological method. In 1990, the nucleic acid sequence of 16s rDNA was analyzed, and it shows

the phylogenetic relationship of Mycobacteria well. Until now, various methods of identifying Mycobacterial species by using the 16S rDNA have been developed and studied (Comparative sequence analysis, Probe hybridization, and Polymerization chain reaction-restriction fragment length polymorphism).

Identifying methods of Mycobacterial species by using dnaJ and 23S rDNA as alternative chronometers were developed in 1994. However, dnaJ and 23S rDNA have problems in phylogenetic relationship determination and conservation of nucleic acid sequences, and thus the methods were not used for target genes (Victor TC, Jordaan AM, Van Schalkwyk EJ, Coetzee GJ, Van Helden PD. Strain-specific variation in the dnaJ gene of mycobacteria. J Med Microbiol. 44(5):332-339, 1996). In 1993, Telenti A et al. reported that a method for the identification of mycobacteria at the species level was developed by using polymerase chain reaction (PCR)-Restriction Enzyme Length Polymorphism of a gene fragment of hsp 65. The method involves steps of amplifying an hsp 65 gene fragment by PCR and restriction enzyme analysis of PCR products of hsp 65 with two restriction enzymes, BstEII and HaeIII, and 29 species and subspecies were differentiated by PCR-restriction enzyme pattern analysis. (Telenti A, Marchesi F, Balz M, Bally F, Bottger EC, Bodmer T. "Rapid identification of mycobacteria to the species level by polymerase chain reaction and restriction enzyme analysis," J. Clin. Microbiol. 31(2):175-8. 1993).

However, the above methods for identifying mycobacteria are

disadvantageous in that the procedure involves various restriction enzymes and is expensive. In addition, the gene fragment must be differentiated to a 10 bp fragment due to the small size of the restriction enzyme fragment in the case of Hae III. Also, mycobacterial species must be identified accurately
5 on the basis of a known restriction fragment database of each species, or they must be analyzed by comparing the electrophoresis patterns of the subject strain and its reference strain.

SUMMARY OF THE INVENTION

10 To resolve the above problems, an object of the present invention is to provide a pair of primers for amplifying the hsp65 gene of mycobacteria.

Another object of the present invention is to provide a polynucleotide of the hsp65 gene fragment that is amplified with the primers.

It is yet another object of the present invention to provide a probe or
15 a probe set for detecting or identifying mycobacterial species comprising at least a gene fragment of the hsp 65 gene of reference mycobacterial species.

It is a still another object of the present invention to provide a simple and accurate method for the detection or identification of mycobacterial species.

20 It is a further object of the present invention to provide a method for the identification of mycobacterial species comprising the steps of:

(1) amplifying an hsp 65 gene fragment of mycobacteria of interest with primers for specifically amplifying the hsp 65 gene of mycobacteria;

(2) analyzing the nucleotide sequence of the amplified hsp 65 gene fragment; and

(3) comparing the nucleotide sequence of the amplified hsp 65 gene fragment obtained in step (2) with a 604-bp hsp 65 gene fragment of a
5 reference mycobacterial species.

It is a further object to provide a method for the detection or identification of mycobacterial species comprising the steps of:

(1) amplifying an hsp 65 gene fragment of mycobacteria of interest with primers for specifically amplifying the hsp 65 gene of mycobacteria; and

10 (2) hybridizing the amplified hsp 65 gene fragment with a probe set comprising at least a probe of the hsp 65 gene fragment.

It is a further object to provide a method for the identification of mycobacterial species comprising the steps of amplifying an hsp 65 gene fragment of mycobacteria of interest with a pair of primers for specifically
15 amplifying the hsp 65 gene of mycobacteria, and analyzing according to the Restriction Fragment Length Polymorphism (RFLP) analysis using the restriction enzyme recognition site in the amplified hsp 65 gene fragment.

It is a further object to provide a kit useful for the diagnosis or identification of mycobacterial species comprising a pair of primers for
20 amplifying the hsp 65 gene of mycobacteria, and a restriction enzyme recognizing the restriction enzyme recognition site which is located in the amplified hsp 65 gene fragment.

It is a further object to provide a kit useful for the diagnosis or

identification of mycobacterial species comprising an amplifying means comprising a pair of primers for specifically amplifying the 604-bp hsp 65 gene fragment of mycobacteria, a hybridization means comprising a probe set including at least a 604-bp hsp 65 gene fragment, and a labeling means
5 for detecting the hybridized product.

BRIEF DESCRIPTION OF THE DRAWINGS

The above and the other objects and features of the present invention will become apparent from the following description given in
10 conjunction with the accompanying drawings, in which:

Fig. 1 shows the hsp 65 gene fragment and the primers of the present invention;

Fig. 2 is a photograph of electrophoresis showing the amplified product of mycobacterial DNA, wherein panel A shows a result obtained from
15 analysis of the amplified gene fragment of reference strains, and panel B shows a result for the amplified gene fragments of mycobacteria in a clinical sample;

Fig. 3 is a photograph of agarose gel electrophoresis showing the hsp 65 gene fragment of a reference mycobacterial species that was
20 amplified and then treated with Xho I;

Fig. 4 is a summarized diagram showing a result of PCR-RFLP of hsp 65 gene fragments of reference strains of mycobacteria;

Fig. 5 is a photograph of agarose gel electrophoresis where hsp

65hsp 65 gene fragments of mycobacteria in a clinical sample were amplified and then treated with XhoI;

Fig. 6 shows phylogenetic relationships of 50 reference mycobacterial species obtained in Example 7; and

5 Figs. 7a to 7d shows the results of the identification of mycobacteria in a clinical sample according to a comparative sequence analysis.

DETAILED DESCRIPTION AND THE PREFERRED **EMBODIMENTS**

10 The present invention relates to a pair of primers specific to mycobacteria, and more specifically to a pair of primers specifically amplifying an hsp 65 gene fragment of mycobacteria, an hsp 65 gene fragment, and a method for the identification of mycobacteria with the same.

Considering the problems in conventional identification methods and
15 the taxonomy of mycobacteria, the inventors provide PCR primers that can amplify *M. tuberculosis* and non-tuberculosis mycobacteria, an hsp 65 gene fragment as a chronometer molecule which exists in all mycobacteria, and a method for the identification of mycobacteria by using the primers and hsp
20 65 gene fragments. By using the restriction fragment of the amplified product of hsp 65 genes with treatment of Xho I, it is possible to differentiate *M. tuberculosis* and non-tuberculosis mycobacteria, and to differentiate non-tuberculosis mycobacteria.

In order to obtain a pair of primers that preferably amplify the hsp 65

gene of mycobacteria, the inventors prepared the primers on the basis of the hsp 65 gene of *M. tuberculosis* (GenBank No. M15467), *M. avium* (GenBank No. AF281650) of which 1623-bp full sequences of the hsp 65 gene were analyzed, and *T. paurometabola* (GenBank No. AF352578) which is phylogenetically closer to mycobacteria. The forward primer comprises 20 nucleotides located at the 163rd position to the 182nd position of the hsp 65 gene sequence of the three mycobacteria, and the backward primer comprises 20 nucleotides located at the 787th position to the 806th position. In addition, the modified primers or polynucleotides comprising the primers can be used for amplifying 644-bp hsp 65 gene fragments of mycobacteria. The primer region of the hsp 65 gene is adopted from the region of *M. tuberculosis* and *M. avium* which belong to genus mycobacteria and *Tsukamurella paurometabola*. Preferably, the forward primer is 5'-ATCGCCAAGGAGATCGAGCT-3', which is called HSPF 3 and is shown in SEQ ID NO: 55. The backward primer is 5'-AAGGTGCCGCGGATCTTGTT-3', which is called HSPR 4 and is shown in SEQ ID NO: 56. The positions of the hsp 65 gene fragment and the primers are schematically indicated in Fig. 1.

The present invention provides polynucleotides of hsp 65 gene fragments used for detecting or identifying mycobacterial species. In addition, the present invention provides polynucleotide sets comprising at least a polynucleotide selected from the group consisting of hsp 65 gene fragments or complementary sequences thereto.

The chronometer molecule used for the identification of mycobacterial species in the present invention is the 644-bp gene fragment located at the 163rd position to the 806th position of a 1623-bp hsp 65 gene of *M. tuberculosis*. The 644-bp gene fragment is substantially a 604-bp fragment because the 40-bp primer sequence is excluded. As a result of a Genbank database search, it was found that all 604-bp gene fragments of hsp 65 of 54 kinds of reference mycobacterial species are novel.

To establish the database for detecting and identifying the mycobacteria, the reference strains as shown in Table 1 were employed. 50 reference strains included 47 reference strains from the American Type Culture Collection (ATCC), a reference strain of *M. leprae* (Thai 53 strains) from Hanssen's disease center of the Catholic University of Korea, and 2 reference strains (type II, III) of *M. kansasii* from V. Vincent. In addition, hsp 65 gene fragments of 3 reference strains of *Tsukamurella* from the German Collection of Microorganisms and Cell Cultures, and a reference strain of *Nocardia* from ATCC were analyzed (Table 1).

[TABLE 1] Reference strains of the present invention

Reference strains of mycobacteria					
No	species	source	No	Species	Source
1	<i>M. abscessus</i>	CAP97E-03	26	<i>M. kansasii</i> Type III	V. Vincent
2	<i>M. africanum</i>	ATCC 25420	27	<i>M. leprae</i>	Thai 53
3	<i>M. asiaticum</i>	ATCC 25276	28	<i>M. malmoeense</i>	ATCC 29571

4	<i>M. aichiense</i>	ATCC 27280	29	<i>M. marinum</i>	ATCC 927
5	<i>M. avium</i>	ATCC 25291	30	<i>M. mucogenicum</i>	ATCC 49650
6	<i>M. bovis</i>	ATCC 19210	31	<i>M. neoaurum</i>	ATCC 25795
7	<i>M. bovis BCG</i>	French strain	32	<i>M. nonchromogenicum</i>	ATCC 19530
8	<i>M. celatum</i> Type I	ATCC 51131	33	<i>M. paratuberculosis</i>	ATCC 19698
9	<i>M. celatum</i> Type II	ATCC 51130	34	<i>M. phlei</i>	ATCC 11758
10	<i>M. chelonae</i>	ATCC 35749	35	<i>M. peregrinum</i>	ATCC 14467
11	<i>M. chitae</i>	ATCC 19627	36	<i>M. scrofulaceum</i>	ATCC 19981
12	<i>M. microti</i>	ATCC 19422	37	<i>M. senegalense</i>	ATCC 35796
13	<i>M. flavescens</i>	ATCC 14474	38	<i>M. shimoidei</i>	ATCC 27962
14	<i>M. fortuitum</i> 6841	ATCC 6841	39	<i>M. simiae</i>	ATCC 25275
15	<i>M. fortuitum</i> 49403	ATCC 49403	40	<i>M. smegmatis</i>	ATCC 19420
16	<i>M. fortuitum</i> 49404	ATCC 49404	41	<i>M. szulgai</i>	ATCC 35799
17	<i>M. gastri</i>	ATCC 15754	42	<i>M. terrae</i>	ATCC 15755
18	<i>M. genavense</i>	ATCC 51233	43	<i>M. thermoresitibile</i>	ATCC 19527

19	<i>M. gordonae</i>	ATCC 14470	44	<i>M. triviale</i>	ATCC 23292
20	<i>M. haemophilum</i>	ATCC 29548	45	<i>M. tuberculosis</i>	ATCC 27294
21	<i>M. interjectum</i>	ATCC 51457	46	<i>M. ulcerans</i>	ATCC 19423
22	<i>M. intermedium</i>	ATCC 51848	47	<i>M. vaccae</i>	ATCC 15483
23	<i>M. intracellulare</i>	ATCC 13950	48	<i>M. wolinskyi</i>	ATCC 700010
24	<i>M. kansasii</i> Type I	ATCC 12478	49	<i>M. parafortuitum</i>	ATCC 19686
25	<i>M. kansasii</i> Type II	V. Vincent	50	<i>M. farcinogenes</i>	ATCC 35753
Reference strain of bacteria other than mycobacteria					
1	<i>T. paurometabola</i>	DSM 20162	2	<i>T. tyrosinosolvens</i>	DSM 44234
3	<i>T. pulmonis</i>	DSM 44142	4	<i>N. carnea</i>	ATCC 6847

For detecting and identifying Mycobacterial species, the present invention provides 604-bp hsp 65 gene fragments as a new chronometer molecule, instead of 16S rDNA. The chronometer molecules must satisfy the following requirements in order to reflect the phylogenetic relationship. Firstly, the target gene must be essential for the functions and be highly conserved in all organisms. Secondly, the target gene must not mutate by lateral transfer based on selection pressure between species. Thirdly, the target gene must have interspecies variation and intraspecies conservation, which suitably reflects a phylogenetic relationship. The hsp 65 gene fragment of the

present invention suitably satisfies the requirements of the chronometer molecule.

The nucleotide sequences of reference strains were analyzed according to a direct sequence analysis method, and compared with each other by multi-alignment. As a result, it was found that strains other than five kinds of TB complex including *M. africanum*, *M. bovis*, *M. bovis BCG*, *M. microti*, and *M. tuberculosis* (54 reference strains) have different nucleotide sequences, namely interspecies variation. In a previous report, the five kinds of mycobacteria belonging to the TB complex had the same nucleotide sequence analyzed according to another analyzing method using the 16S rDNA or *rpoB* gene fragment, and it was found that the mycobacterial species belonged to the same species. The result showed that the hsp 65 gene fragment of the present invention satisfied the interspecies variation of nucleotide sequences. Secondly, all 54 reference strains used in the experiment have 604-bp coding sequences without insertion and deletion, when they are multi-aligned. That is, there is no gap on multi-aligned sequences. 16S rDNA has a gap at a high frequency on multi-alignment of the nucleotide sequence. It is known that because an aligned gene corresponding to the gap must be eliminated in performing multi-alignment, the gap causes an error in establishing the phylogenetic tree at a high rate. Therefore, the identification method using the hsp 65 gene of the present invention provides significant advantages.

In order to investigate whether the 604-bp hsp 65 gene fragment of

the present invention can be suitable for use as a chronometer molecule, a phylogenetic tree was constructed by the nucleotide sequence of 604-bp hsp 65 gene fragments of various mycobacteria. In addition, the mycobacteria identified according to the other conventional method were analyzed by the
5 identification method of the present invention using the hsp 65 gene fragment. As a result, the present invention accurately identified the mycobacteria.

The phylogenetic tree of the reference strains of the present invention showed the natural relationships of the mycobacteria. That is, the
10 result confirmed that 50 reference strains of TB complex formed a large group excluding *T. paurometabola* as an outgroup (Fig. 6). Also, slow-growing mycobacteria and fast-growing mycobacteria formed different groups. *M. tuberculosis* and *M. leprae* of pathogenic mycobacteria formed the same branch of the phylogenetic tree. MOTT were isolated frequently. *M.*
15 *avium* and *M. intracellulare*, showing quite similar biochemical characteristics, formed the same branch. The results showed general characteristics of mycobacteria. *M. kansasii* and *M. gastri* have 100% sequence homology, and thus cannot be differentiated according to the conventional identification method using 16S rDNA, but they are differentiated according to the present
20 invention. Moreover, the subspecies of *M. kansasii* can be differentiated (namely, the hsp 65 gene fragments of *M. kansasii* Type I, II, and III have different nucleotide sequences). The results of the present invention show the phylogenetic relationships of mycobacteria. That is, the slow-growing

mycobacteria and fast-growing bacteria form different branches of the phylogenetic tree, and *M. tuberculosis* and *M. lepre* form the same branch.

The mycobacterial species can be identified according to the identifying method of mycobacteria, such as comparative sequence analysis, probe hybridization, and PCR-RFLP, using the polynucleotide of the present invention. The comparative sequence analysis, probe hybridization, and PCR-RFLP can be carried out according to the method which has been known to a skilled person in the art. For example, a method for identifying the mycobacteria with 16s rDNA can be applied for the identification method of the present invention.

In one aspect, the present invention provides a method for the identification of mycobacteria by using PCR-RFLP (also called PRA). The method comprises the steps of amplifying hsp 65 gene fragments of mycobacteria with primers specific to mycobacteria, preferably primers as shown in SEQ ID NOs: 55 and 56, and analyzing the amplified product according to the RFLP analysis by using the restriction enzyme recognizing the target site located in the amplified product. The identification method is simple, economical, and specific to the mycobacteria.

The general PRA method comprising the steps of 1) DNA extraction, 2) PCR amplification, 3) confirmation of the amplified product, 4) digestion with a restriction enzyme, 5) analysis of restriction fragment, and 6) visualization by image capture systems. The restriction enzymes applicable for this invention include all the restriction enzymes that can recognize the

site specifically existing in a 644-bp fragment, preferably Xho I.

In addition, any method for differentiating the restriction fragment on the basis of its size, preferably electrophoresis, and more preferably agarose gel electrophoresis or polyacrylamide gel electrophoresis, can be applied for
5 analyzing the restriction fragment.

In comparison with the conventional method for identifying mycobacteria using the hsp 65 gene, the method of the present invention is simple and economical. The conventional method uses a 439-bp fragment of hsp 65 gene as a target gene, and two kinds of restriction enzymes, Hae II
10 and BstE II. As described above, the identification method of the present invention uses one restriction enzyme, such as Xho I, and it is advantageous in time and cost. In addition, the 439-bp fragment of the conventional method is shorter than 644-bp of the present invention. By treating with Hae III and recognizing four nucleotides in the conventional method, many fragments
15 are produced so that the small fragments, such as a 10-bp fragment, must be separated. Thus, in order to accurately identify mycobacteria in the conventional method, it is necessary to use the restriction fragment database of reference strains, or to analyze the mycobacteria of interest together with putative reference species according to restriction enzyme treatment and
20 electrophoresis. In the preferred embodiment of the present invention, the identifying method of mycobacteria uses Xho-I recognizing six (6) nucleotides as target sites, thereby making it perform more gel electrophoresis. However, the identification method is accurate and simple.

- The present invention provides a new system where a 644-bp hsp 65 gene fragment of mycobacteria is amplified with primers specifically for amplifying the 644-bp hsp 65 gene fragment of mycobacteria, and it is treated with Xho-I to differentiate and identify the mycobacterial species.
- 5 Only a process of PRA makes it possible to differentiate the MOTT into 3 groups, as well as *M. tuberculosis*. That is, the treatment of the amplified product with a restriction enzyme produces only a 644-bp gene fragment in fast-growing mycobacteria, thereby differentiating it from the slow-growing mycobacteria. *M. avium* complex (for examples, *M. avium* and *M.*
- 10 *intracellulare*) which belongs to slow-growing mycobacteria and is isolated most frequently in clinical samples produces three kinds of restriction fragments, 391-bp, 169-bp, and 84-bp, thereby differentiating them from other groups including *M. kansasii* producing two kinds of restriction fragments, 391-bp and 253-bp.
- 15 Among the genus *Mycobacterium* that includes about 70 species, about 10 strains including *M. tuberculosis*, *M. avium* complex, *M. kansasii*, *M. szulgai*, *M. gordonae*, *M. fortuitum*, and *M. chelonae* cover 90% of isolates in a clinical sample, and thus they can be effectively identified according to the identification method of the present invention.
- 20 In another aspect of the present invention, a TB complex can be differentiated from MOTT by treating the amplified hsp 65 gene fragment with Xho I, and analyzing it according to RFLP. In addition, the TB complex can be differentiated based on the restriction fragment of the amplified 644-

bp hsp 65 fragments of 391-bp, 150-bp, and 103-bp.

In a further aspect of the present invention, a 644-bp hsp 65 gene fragment of fast-growing mycobacteria is not cleaved by a restriction enzyme, Xho I. The fast growing mycobacteria can be differentiated depending on
5 whether the amplified product can be cleaved by the restriction enzyme or not. Thus, the present invention provides a method for differentiating fast-growing mycobacteria among MOTT. When 391-bp, 169-bp, and 84-bp restriction fragments are produced by the treatment of Xho I and RFLP analysis of mycobacteria, the mycobacteria can be identified as species
10 including *M. avium*, *M. intracellulare*, *M. celatum*, *M. shimoidei*, and *M. szulgai*.

In the case that the treatment of Xho I and the RFLP analysis produces 391-bp and 253-bp restriction fragments, the mycobacterial species are identified as species including *M. gastri*, *M. genavense*, *M.*
15 *gordonae*, *M. haemophilum*, *M. kansasii*, *M. malmoense*, *M. marinum*, *M. scrofulaceum*, *M. simiae*, and *M. ulcerans*.

The differentiation results are summarized, depending on the size pattern of restriction fragments obtained in the present invention, in Fig. 4.

The present invention also relates to a kit for differentiating or
20 diagnosing mycobacterial species comprising Xho I and primers specific to the hsp 65 gene of mycobacterial species, preferably primers as shown in SEQ ID NOs: 55 and 56, wherein the DNA of mycobacterial species in a sample is amplified with the primers to produce the hsp 65 gene fragment,

and the mycobacterial species are differentiated depending on the restriction fragments obtained according to RFLP. The kit further comprises a PCR amplification kit and a RFLP kit. Any kit that has been known for the use and is commercially available is applicable to the present invention.

5 In another aspect, the present invention relates to a method for detecting and identifying the mycobacterial species, comprising the steps of (1) amplifying 604-bp hsp 65 gene fragments of mycobacterial species of interest with a primer that can specifically amplify hsp 65 gene fragments, (2) hybridizing the amplified product with a probe set comprising at least a 604-
10 bp hsp 65 gene fragment selected from the group consisting of the polynucleotide of a 604-bp hsp 65 gene fragment of mycobacterial species. In the embodiment of the method, the hsp 65 gene fragment of mycobacterial species of interest can be amplified according to general amplification methods of nucleotides such as PCR, LCR (ligase chain
15 reaction), NASBA, etc. The amplified product can be hybridized with 604-bp gene fragment(s) or 644-bp gene fragment(s) of the reference species, preferably a probe or probe set comprising a fragment selected from the group consisting of polynucleotides as shown in SEQ ID NO: 1 to SEQ ID NO: 54, and the polynucleotides complementary thereto. The hybridization
20 step can be carried out according to a general hybridization method. For example, the hybridization can be performed on a solid surface, or it can be carried out by using a microarray including the probe immobilized thereon.

The present invention provides an identification or diagnosis kit

comprising (1) a means for amplification including a pair of primers specific to an hsp 65 gene of mycobacterial species; (2) a means for hybridization comprising a 604-bp or 644-bp gene fragment of hsp 65 of mycobacterial species, preferably a probe or probe set comprising a gene fragment selected from the group consisting of polynucleotides as shown in SEQ ID NO: 1 to SEQ ID NO: 54 and polynucleotides complementary thereto; and (3) a labeling means for detecting the hybridized product. The labeling means can include all the labeling means that can be generally used for detection of hybridized DNA, for example Cy5, biotin-binding compounds, Cy3, EDANS(5-(2'-aminoethyl)amino-1-naphthalene sulfate), tetramethylrhodamine (TMR), tetramethylrhodamine isocyanate (TMRITC), x-rhodamine, and Texas red.

The present invention provides a method for identification of mycobacterial species by using comparative sequence analysis. The database of hsp 65 gene fragments as described above can be applicable to the method. The present invention provides a method for identification of mycobacterial species by using 604-bp hsp 65 gene fragments. More specifically, the method comprises the steps of:

- (1) amplifying hsp 65 gene fragments of mycobacteria of interest with primers for specifically amplifying the hsp 65 gene of mycobacteria;
- (2) analyzing the nucleotide sequence of the amplified hsp 65 gene fragment; and
- (3) comparing the nucleotide sequence of the amplified hsp 65 gene

fragment obtained in step (2) with a 604-bp hsp 65 gene fragment of a reference strain of mycobacteria.

Preferably, step (3) can be carried out by multi-aligning the 604-bp hsp 65 gene fragment of mycobacterial species of interest with a polynucleotide set comprising at least an hsp 65 604-bp polynucleotide of reference strains of mycobacteria to infer a phylogenetic tree. According to the comparative sequence analysis, a database of 604-bp hsp 65 gene fragments is established by amplifying the hsp 65 gene fragment of reference species of mycobacteria with the primers specific to mycobacteria, preferably primers as shown in SEQ ID NO: 55 and SEQ ID NO: 5, and analyzing the nucleotide sequence of the amplified product. In the example of the present invention, the database of 604-bp fragments of 54 reference strains except for the primer sequence is established by analyzing the nucleotide sequences of the 604-bp fragments, and through multi-alignment.

15 The 604-bp fragments of reference strains obtained in the present invention are shown in SEQ ID NO: 1 to SEQ ID NO: 54. The mycobacterial species of interest can be identified according to comparative sequence analysis by using the database.

As the hsp 65 gene fragments of the mycobacterial species of interest are different from those of the reference species, mycobacterial species of interest can be identified based on the criterion of nucleotide sequence homology of hsp 65 genes of reference species. Because a mycobacterial species has a different range of sequence homology,

20

mycobacterial species can be identified based on the specific range of the sequence homology thereof. For example, *M. gordonae* has a wide range of sequence homology, but *M. tuberculosis* has a narrow range. In addition, mycobacterial species can be identified by multi-aligning the nucleotide
5 sequence of 604-bp hsp 65 gene fragments with those of reference species to infer a phylogenetic relationship.

To confirm that the database including 604-bp hsp 65 gene fragments of 50 reference strains of mycobacterial species can be useful for identifying the mycobacterial species in a clinical sample, the identification
10 method of the present invention was applied for 38 strains of mycobacteria obtained from the Korean Institute of Tuberculosis of the Korean National Tuberculosis Association, which had already been identified by using biochemical identification methods such as pigmentation on solid media; optimal growth temperature; degree of growth on media including catalase,
15 iron, and p-nitrobenzoic acid; hydrolysis of tween 80; a Tellulite reduction test; degree of growth on media including 5% NaCl; production of Niacin; Nitrate reduction test; and production of Urease according to a blind test. The test results are shown in Table 2. In the table, the strain item indicates the number offered by the Korean Institute of Tuberculosis at random. The
20 biochemical method item is a result of identification by the Korean Institute of Tuberculosis, and the item hsp 65 gene analysis method is a result of the present invention.

[Table 2] Identification result for clinical isolates

No.	strain	Biochemical method	hsp 65 gene analysis method
1	KIT 77009	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
2	KIT 77710	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
3	KIT 77712	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
4	KIT 77714	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
5	KIT 77719	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
6	KIT 77720	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
7	KIT 77721	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
8	KIT 77722	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
9	KIT 77723	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
10	KIT 77725	<i>M. tuberculosis</i>	<i>M. tuberculosis</i>
11	KIT 41105	<i>M. avium</i> complex	<i>M. intracellulare</i>
12	KIT 41110	<i>M. avium</i> complex	<i>M. avium</i>
13	KIT 41111	<i>M. avium</i> complex	<i>M. intracellulare</i>
14	KIT 41115	<i>M. avium</i> complex	<i>M. intracellulare</i>

15	KIT 30101	<i>M. scrofulaceum</i>	<i>M. scrofulaceum</i>
16	KIT 30102	<i>M. scrofulaceum</i>	<i>M. scrofulaceum</i>
17	KIT 20118	<i>M. kansasii</i>	<i>M. kansasii</i> Type I
18	KIT 20119	<i>M. kansasii</i>	<i>M. kansasii</i> Type I
19	KIT 20120	<i>M. kansasii</i>	<i>M. kansasii</i> Type I
20	KIT 47101	<i>M. terrae</i> complex	<i>M. nonchromogenicum</i>
21	KIT 47102	<i>M. terrae</i> complex	<i>M. nonchromogenicum</i>
22	KIT 47103	<i>M. terrae</i> complex	<i>M. nonchromogenicum</i>
23	KIT 47104	<i>M. terrae</i> complex	<i>M. nonchromogenicum</i>
24	KIT 32101	<i>M. gordonae</i>	<i>M. gordonae</i>
25	KIT 32104	<i>M. gordonae</i>	<i>M. gordonae</i>
26	KIT 32105	<i>M. gordonae</i>	<i>M. gordonae</i>
27	KIT 32106	<i>M. gordonae</i>	<i>M. gordonae</i>
28	KIT 31102	<i>M. szulgai</i>	<i>M. szulgai</i>
29	KIT 31103	<i>M. szulgai</i>	<i>M. szulgai</i>

30	KIT 31106	<i>M. szulgai</i>	<i>M. szulgai</i>
31	KIT 31107	<i>M. szulgai</i>	<i>M. szulgai</i>
32	KIT 21101	<i>M. marinum</i>	<i>M. marinum</i>
33	KIT 60108	<i>M. fortuitum</i> complex	<i>M. fortuitum</i> 6841
34	KIT 60109	<i>M. fortuitum</i> complex	<i>M. fortuitum</i> 6841
35	KIT 60110	<i>M. fortuitum</i> complex	<i>M. fortuitum</i> 6841
36	KIT 60111	<i>M. fortuitum</i> complex	<i>M. fortuitum</i> 6841
37	KIT 61104	<i>M. chelonae</i> complex	<i>M. abscessus</i>
38	KIT 61105	<i>M. chelonae</i> complex	<i>M. abscessus</i>

The nucleotide sequences of 38 mycobacteria obtained from the clinical sample were analyzed and then multi-aligned with the database of reference strains to infer the phylogenetic tree. From the results, all 38 strains were identified to the species level with 100% sensitivity and specificity (Table 2 and Figs. 7a to 7d). The results are specifically described below.

A. Identification of *M. tuberculosis*

M. tuberculosis is the most pathogenic and important species in public health. The results of the identification of *M. tuberculosis* by using the

database of the reference species of mycobacteria of the present invention confirmed that all twenty (20) *M. tuberculosis* were identified (Table 2 and Fig. 7c), and showed that 604-bp hsp 65 gene fragments of 20 strains have 100% sequence homology with a 604-bp fragment of *M. tuberculosis* ATCC 27284 reference strain. The 16s rDNA and rpoB gene used as a target gene are involved in resistance to streptomycin and to rifampin, respectively. The target genes in mycobacteria with a resistance to antituberculosis drugs can be mutated. However, unlike 16s rDNA and rpoB, the hsp 65 gene is not related to resistance to antibiotics, and thus it does not mutate. Therefore, the 604-bp hsp 65 gene is stable with respect to the selection pressure of antituberculosis drugs in comparison with other target genes.

B. Identification of *M. avium* complex

The identification method was applied to 4 strains of *M. avium* complex which are the most commonly isolated in MOTT. As a result, the strains were identified to the species level as 3 *M. intracellulares* and a strain of *M. avium*. According to the biochemical identification method, it is not possible to differentiate *M. intracellulare* and *M. avium* because they have the same biochemical characteristics. In comparing the nucleotide sequences of 1 *M. avium* (KIT 41110) and *M. avium* ATCC 25281, they have 99.5% nucleotide sequence homology with 3 different nucleotides. When the nucleotide sequences of the 3 *M. intracellulares* (KIT 41105, 41111, and 51115) are compared with that of *M. Intracellulare* ATCC 13850, they show 99.0-99.8% sequence homology. Those results are consistent with the fact

that *M. intracellulare* includes various genotypes, namely interspecies heterogeneity (Devallois A, Picardeau M, Paramasivan CN, Vincent V, Rastogi N: Molecular characterization of *Mycobacterium avium* complex isolates giving discordant results in AccuProbe tests by PCR-restriction enzyme analysis, 16s rRNA sequencing, and DT1-DT6 PCR. *J Clin Microbiol* 1997 35: 2767-2772).

C. Identification of *M. scrofulaceum*.

2 strains (KIT 30101, 30102) were identified as *M. scrofulaceum* (Fig.7b), and they have 99.8-100% nucleotide sequence homology with *M. scrofulaceum* ATCC 19981.

D. Identification of *M. kansasii*

M. kansasii is the most pathogenic in MOTT, and it is in second position in isolation frequency after *M. avium* complex. 16s rDNA of *M. kansasii* showed 100% nucleotide sequence homology with that of non-pathogenic *M. gastri*, thereby making it difficult to differentiate them. In addition, *M. kansasii* consists of at least 5 subspecies where type II and type III are reported to be separated from clinical material. 3 strains are identified as *M. kansasii* by using the database of 604-bp hsp 65 gene fragments of reference strains, which are consistent with results of the biochemical identification method. The method for identifying the mycobacterial species by using the database has characteristics such that *M. kansasii* can be differentiated from *M. gastri*, and subspecies of *M. kansasii* can be differentiated. The result confirmed that 3 strains of *M. kansasii* (KIT 20118,

20119, 20120) have 100% nucleotide sequence homology, and they are identified as *M. kansasii* Type I ATCC 12478 (Fig. 4c).

E. Identification of *M. gordonae*, *M. szulgai*, *M. marinum*, and *M. terrae* complex

5 As a result of identifying the clinically separated mycobacterial strains with the database of the present invention, 4 strains (KIT 32101, 32104, 32105, and 32106) were found to be *M. gordonae* (Fig. 7a, and Table 2). When comparing the nucleotide sequences of the 604-bp hsp 65 gene fragments of the 4 strains, they have 99.2-99.8% sequence homology with
10 each other, but they have 95.9-96.3% sequence homology with *M. gordonae* ATCC 14470, which indicates a considerably low sequence homology between *M. gordonae* species. The result is consistent with the report that *M. gordonae* has intraspecies heterogeneity (Abed Y, Bollet C, de Micco P. Identification and strain differentiation of Mycobacterium species on the basis
15 of DNA 16S-23S spacer region polymorphism. Res Microbiol. 1995 146(5): 405-13). That is, 4 isolates obtained from the same region have high sequence homology with one another, but low sequence homology with reference strains obtained from different regions.

 As a result of identification of mycobacterial species with the
20 database of reference species, 4 strains (KIT 31102, 31103, 31106, and 31107) were identified as *M. szulgai*, which is consistent with that of the biochemical identification method (Fig. 7a, and Table 2). The nucleotide sequences of the 4 strains have 99.5-100% nucleotide sequence homology

with *M. szulgai* ATCC 35799.

As a result of identification of mycobacterial species with the database of reference species, 1 strain was identified as *M. marinum*, which is consistent with that of the biochemical identification method (Fig. 7a and Table 2). The nucleotide sequence of the strain has 99.3% nucleotide sequence homology with *M. marinum* ATCC 927(//).

M. terrae complex does not generally cause disease in humans, and it includes 3 reference strains of the present invention (*M. terrae*, *M. triviale*, *M. nonchromogenicum*), and various mycobacterial species which are not classified. As a result of identification of mycobacterial species with the database of reference species, 4 strains were identified as *M. nonchromogenicum* among the *M. terrae* complex, which is consistent with the previous report using the biochemical identification method. When comparing the 4 strains with the reference strain, the nucleotide sequences of the strains had 95.0-100% nucleotide sequence homology with *M. nonchromogenicum* ATCC 19530. The result concurs with the report that *M. terrae* complex has intraspecies heterogeneity.

F. Identification of fast-growing mycobacteria (*M. fortuitum* complex and *M. chelonae* complex)

As a result of identification of mycobacterial species with the database of reference species, 2 strains (KIT 61104, 61105) were identified as *M. abscessus* of *M. chelonae* complex. The result is consistent with that of the biochemical identification method. However, the identification method

of the present invention resolves the problem of the conventional biochemical method that *M. chelonae* and *M. abscessus* cannot be differentiated. The nucleotide sequences of the hsp 65 gene fragments of the strains have 98.4-99.5% nucleotide sequence homology with *M.*
5 *abscessus* CAP97E-03.

According to the identification method of the present invention, 4 strains are identified as *M. fortuitum*, which is consistent with the result of the biochemical identification method. *M. fortuitum* complex covers various mycobacterial species, and includes *M. fortuitum* ATCC 6841, *M. fortuitum*
10 ATCC 49403, *M. fortuitum* ATCC 49404, and *M. peregrinum* as reference species. 4 clinically isolated strains were identified as *M. fortuitum* ATCC 6841. When the 4 strains are compared with *M. fortuitum* ATCC 6841, they have 99.4 ~ 100% nucleotide sequence homology.

The present invention is further specifically illustrated in the following
15 examples, which should not be taken to limit the scope of the invention.

EXAMPLE 1

DNA Isolation from reference strains and clinically isolated strains

20 1-1: Selection of subject strain

As shown in Table 1, The hsp 65 gene fragments of 50 reference strains were sequenced, including 47 reference strains from the American Type Culture Collection (ATCC), a reference strain of *M. leprae* (Thai 53

strain) from Hanssen's disease center of the Catholic university of Korea, and 2 reference strains (type II, III) of *M. kansasii* from V. Vincent. In addition, 3 reference strains of *Tsukamurella* from the German Collection of Microorganisms and Cell Cultures and a reference strain of *Nocardia* from 5 ATCC were selected.

As shown in Tables 2 and 4, the clinically isolated strains are strains that were previously identified according to the biochemical identification method.

1-2: DNA Isolation

10 The genomic DNA of reference strains and clinically isolated strains was extracted according to the Bead Beater Phenol (BB/P) extraction method. The culture of each mycobacteria was suspended with TEN buffer (10 mM Tris-HCl, 1 mM EDTA, 100 mM NaCl: pH 8.0) placed in a 2.0 ml screw-cap microcentrifuge tube filled with 100 μl (packed volume) of beads 15 (glass beads, 0.1 mm diameter; Biospec Products, Bartlesville, Okla., U.S.A.) and 100 μl phenol:chloroform:isopropylalcohol solution (50:49:1, v/v/v). To disrupt the bacteria, the tube was oscillated with a Mini-Bead beater (Biospec Products, Bartlesville, Okla., U.S.A.) for one minute, and to separate phases, the tube was centrifuged (12,000 rpm, 5 min). After the 20 supernatant (100 μl) was transferred into another clean tube, the tube was centrifuged at 15,000 rpm for 5 min with the addition of 60 μl of isopropyl alcohol. The resultant DNA pellet was washed with 70% ethanol, solubilized with 60 μl of TE buffer (10 mM Tris-HCl, 1 mM EDTA), and used as a

template DNA for the analysis of nucleotide sequence and identification of the mycobacterial species in the following Examples.

EXAMPLE 2

5 **Preparation of primers for amplifying hsp 65 gene fragments**

A forward primer and a backward primer were prepared for specifically amplifying hsp 65 genes of all mycobacterial species. *hsp 65* genes of *M. tuberculosis* (GenBank No. M15467) and *M. avium* (GenBank No. AF281650) of which 1623-bp full sequences were previously analyzed for
10 another purpose, and *T. paurometabola* (GenBank No. AF352578) were used for this example to prepare primers for amplifying hsp 65 genes of all the mycobacteria. The primers were shown in SEQ ID NO: 55 and 56, and positions thereof are indicated in Fig. 1.

Forward primer: HSPF3 (SEQ ID NO: 55)

15 5'-ATCGCCAAGGAGATCGAGCT-3'

Backward primer: HSPR4 (SEQ ID NO: 56)

5'-AAGGTGCCGCGGATCTTGTT-3'

EXAMPLE 3

20 **Amplification of 644-bp hsp 65 gene fragment**

3-1) PCR amplification of *hsp 65* gene

PCR reaction was carried out using AccuPower PCR PreMix (Korea, bioneer) containing 2 U Taq polymerase, 10 mM dNTP, 10 mM Tris-HCl (pH

8.3), and 1.5 mM MgCl₂. 50 ng of each DNA isolated in Example 1, and 20 pmol of each primer prepared in Example 2 were placed in a tube and distilled water was added thereto to a final volume of 20 μ l. PCR was performed at 95 °C for 5 min for a first denaturation followed by 30 cycles of
5 1 min at 95 °C for subsequent denaturation, 45s at 62 °C for annealing, 1 min 30s at 72 °C for extension, and 5 min at 72 °C for final extension (Model 9600 thermocycler, Perkin-Elmer cetus). After PCR reaction, PCR products were electrophoresed on 1% agarose gel to observe a 644-bp fragment.

10 3-2) Separation of PCR product

After electrophoresis on 1% gel, a gel part containing the 644-bp of PCR product was cut and transferred into a new tube in order to separate DNA. DNA isolation and purification were carried out using a Qiaex (Qiagen, Germany) system. 500 μ l of a solution for gel dissolution, QX1, was
15 added to the tube, and the gel and solution were melted at 50 °C for 15 min. Then, 10 μ l of gel beads were mixed therein and held at 50 °C for 15 min. The tube was subjected to a vortex for 10s at intervals of 1 min to equally spread the beads. The tube contents were then washed once with QX1 and twice with QF, dried at 45 °C for 10 min, followed by addition of a TE buffer to
20 obtain 20 μ l of DNA.

After PCR reaction, 1% agarose gel electrophoresis confirmed that a 644- bp fragment was obtained, which is shown in Fig. 2.

Panel A in Fig. 2 indicates amplified DNA products of reference

strains as follows.

Lane M: DNA size marker obtained by treating 174 with *Hae* III;

- | | | |
|----|-------------------------------|-------------------------------|
| | 1: <i>M. tuberculosis</i> , | 2: <i>M. bovis</i> , |
| | 3: <i>M. africanum</i> , | 4: <i>M. avium</i> , |
| 5 | 5: <i>M. intracellulare</i> , | 5: <i>M. scrofulaceum</i> , |
| | 6: <i>M. gordonae</i> , | 7: <i>M. szulgai</i> , |
| | 8: <i>M. marinum</i> , | 9: <i>M. ulcerans</i> , |
| | 10: <i>M. celatum</i> Type I, | 11: <i>M. genavense</i> , |
| | 12: <i>M. malmoense</i> , | 13: <i>M. fortuitum</i> 6841, |
| 10 | 14: <i>M. abscessus</i> , | 15: <i>M. chelonae</i> , |
| | 16: <i>M. peregrinum</i> . | |

Panel B in Fig. 2 indicates amplified DNA products of clinically isolated strains as follows.

lane M: DNA size marker obtained by treating 174 with *Hae*III;

- | | |
|----|---|
| 15 | 1-4: Tbc – clinically isolated strain of <i>M. tuberculosis</i> |
| | 5-7: Mac – clinically isolated strain of <i>M. avium</i> complex; |
| | 8-10: Kac – clinically isolated strain of <i>M. kansasii</i> ; |
| | 11-13: Foc – clinically isolated strain of <i>M. fortuitum</i> ; |
| | 14-16: Chc – clinically isolated strain of <i>M. chelonae</i> . |

20 As shown in Fig. 2, 644-bp hsp 65 gene fragments were obtained from reference strains and clinically isolated strains used in the Example. Therefore, the result suggests that the primers of the present invention could amplify the hsp 65 gene of all the mycobacteria.

EXAMPLE 4

Nucleotides sequence analysis of hsp 65 gene fragment

4-1: Sequence analysis

5 Two strands of 604-bp hsp 65 gene fragments except for 40-bp of primer region which corresponded to the 183rd to 806th positions in hsp 65 of *M. tuberculosis* were sequenced with a forward primer (HSPF3) and a backward primer (HSPR4). The eluted DNA from the gel was used as a template, and automatic sequencing was performed. 1060 ng of the template

10 DNA, 1.2 pmol of each primer, and 2 μl of dye from a BigDye Terminator Cycle Sequencing kit (PE Applied Biosystems) were mixed, and distilled water were added thereto, to a final volume of 10 μl . Reaction was undertaken with a Perkin Elmer Cetus 9600 for 25 cycles of 10 sec at 95 °C, 10 sec at 60 °C, and 4 min at 60 °C. DNA was purified from the reacted

15 sample by an ethanol precipitation method. That is, after 180 μl of distilled water and 10 μl of 3 M sodium acetate were added to the sample to bring the total volume to 200 μl , twice the volume of 100% ethanol was mixed with the mixture and centrifuging was carried out to precipitate DNA. After adding 500 μl of 70% ethanol, centrifuging was carried out at 15,000 rpm

20 for 20 min to wash the DNA. The DNA was recovered with formamide (PE Applied Biosystems). The purified DNA was incubated at 95 °C for 5 min to generate single strand DNA, and the sequence was analyzed with an ABI

3100 system (ABI3100, PE Applied Biosystems) after electrophoresis for 2 hours 30 min. From a search on Genbank, all 604-bp hsp 65 gene fragments of 54 reference strains were found to be novel.

4-2) Alignment of 604-bp hsp 65 gene fragment

5 The nucleotide sequences obtained in the examples were multi-aligned by using the Megalign program of the Dnastar software to construct a database of hsp 65 gene fragments.

 The *rpoB* nucleotide sequence (606-bp) of 54 reference strains analyzed by EXAMPLE 4-2) were multi-aligned by using the Megalign
10 program of the Dnastar software to construct a database of hsp 65 gene fragments.

 For the multiple alignment, 604- bp nucleotides were translated to 301 amino acid residues and the amino acid residues were multiply aligned by a Clustal Method of the Megalign program. The database for identifying
15 the Mycobacteria was constructed using 604 bp nucleotides deduced from the aligned 301 amino acid residues. Sequence homology among nucleotide sequences of reference strains was determined by analyzing multiple aligned database with sequence distance on the Megalign program.

 The nucleotide sequences of 54 reference strains were analyzed by
20 using a direct sequence analysis method, and then multi-aligned. The result confirmed that other reference strains except for TB complex including *M. africanum*, *M. bovis*, *M. vobis* BCG, *M. microti*, and *M. tuberculosis* had different sequences.

4-3) Construction of Phylogenetic tree

The phylogenetic relationship between reference strains was analyzed using a phylogenetic tree constructed by MEGA software (Kumar, S., K. Tamura, and N. Masatoshi. 1993. MEGA: molecular evolutionary genetics analysis, version 1.01. Pennsylvania State University, University Park). The multiple aligned 604-bp polynucleotides from 50 kinds of mycobacterial species were used to construct a Neighbor-joining phylogenetic tree based on the Juke-Cantor distance estimation method and a pairwise deletion method by using a 604-bp polynucleotide of *T. paurometabola* as a outgroup. An analysis of bootstrap was performed through 100 replications. 50 kinds of mycobacteria reference strains made a large group, and fast-growing mycobacteria and slow-growing mycobacteria were formed into different groups from each other. The result reflected the general characteristics of mycobacteria in that pathogenic mycobacteria, *M. tuberculosis*, and *M. leprae* were located in the same branch, and *M. avium* and *M. intracellulare* among MOTT were also formed in the same branch on the phylogenetic tree. In addition, the phylogenetic tree was characterized in that *M. kansasii* could be differentiated from *M. gastri*, which could not be differentiated according to the identification method using 16s rDNA. Also, subspecies of *M. kansasii* can be differentiated (that is, nucleotide sequences of Type I, II, and III of *M. kansasii* are different) (Fig. 6).

EXAMPLE 5**Differentiation of reference strains of mycobacteria by using the PRA**

Xho-I (5'-CTCGAG-3') with 6 nucleotide recognition sites was
 5 determined by analyzing 644-bp (corresponding to the 163rd to the 806th
 position) of hsp 65 gene of *M. tuberculosis* (GenBank No. M15467) and *M.*
avium (GenBank No. AF281650) with the Mapdraw program of Dnastar
 software.

10 10 ul of the 644-pb PCR product in EXAMPLE 3 of 27 kinds of
 reference strains as shown in Table 3 were transferred to a new test tube,
 treated with 1ul (10 units) of Xho-I, and became a final volume of 20 ul with
 addition of 2 ul of 10X buffer and distilled water. The mixture was kept at
 37 °C in a bath for 1 hour. The cleaved mixture was analyzed by 2% agarose
 gel electrophoresis in order to investigate whether the 27 kinds of reference
 15 strains could be identified.

[Table 3] PRA analysis of 27 kinds of reference strains

TB complex		
No.	strain	Source
1	<i>M. africanum</i>	ATCC 25420
2	<i>M. bovis</i>	ATCC 19210
3	<i>M. bovis</i> BCG	French strain
4	<i>M. tuberculosis</i> H37Rv	ATCC 27294
Slow-growing mycobacteria		

5	<i>M. avium</i>	ATCC 25291
6	<i>M. celatum Type I</i>	ATCC 51131
7	<i>M. celatum Type II</i>	ATCC 51130
8	<i>M. gastri</i>	ATCC 15754
9	<i>M. genavense</i>	ATCC 51233
10	<i>M. gordonae</i>	ATCC 14470
11	<i>M. haemophilum</i>	ATCC 29548
12	<i>M. interjectum</i>	ATCC 51457
13	<i>M. intracellulare</i>	ATCC 13950
14	<i>M. kansasii Type I</i>	ATCC 12478
15	<i>M. malmoense</i>	ATCC 29571
16	<i>M. marinum</i>	ATCC 927
17	<i>M. scrofulaceum</i>	ATCC 19981
18	<i>M. shimoidei</i>	ATCC 27962
19	<i>M. simiae</i>	ATCC 25275
20	<i>M. szulgai</i>	ATCC 35799
21	<i>M. ulcerans</i>	ATCC 19423
Rapid-growing mycobacteria		
22	<i>M. abscessus</i>	CAP97E-03
23	<i>M. chelonae</i>	ATCC 35749
24	<i>M. chitae</i>	ATCC 19627
25	<i>M. fortuitum</i> 49403	ATCC 49403
26	<i>M. fortuitum</i> 6841	ATCC 6841
27	<i>M. peregrinum</i>	ATCC 14467

Differentiation of TB complex and MOTT

4 kinds of reference strains (*M. tuberculosis*, *M. bovis*, *M. bovis* BCG, *M. africanum*) which belong to a TB complex of strongly pathogenetic mycobacteria could be differentiated from opportunistic pathogens of MOTT

by use of specific restriction fragments of 391-bp, 150-bp, and 103-bp. The results are shown in Figs. 3 and 4.

Fig. 3 is a photograph of agarose gel electrophoresis of a 644-bp PCR product of an hsp 65 gene fragment of reference strain treated with Xho

5 I.

Lane M: DNA size marker obtained by treating 174 with *Hae* III,

1: <i>M. tuberculosis</i> ,	2: <i>M. bovis</i> ,
3: <i>M. bovis</i> BCG,	4: <i>M. africanum</i> ,
5: <i>M. avium</i> ,	6: <i>M. intracellulare</i> ,
7: <i>M. celatum</i> Type I,	8: <i>M. ulcerans</i> ,
9: <i>M. gordonae</i> ,	10: <i>M. asiaticum</i> ,
11: <i>M. marinum</i> ,	12: <i>M. kansasii</i> ,
13: <i>M. fortuitum</i> 6841,	14: <i>M. abscessus</i> ,
15: <i>M. chelonae</i> ,	16: <i>M. peregrinum</i>

10

15 Lanes 1-4 indicated that *M. tuberculosis*, *M. bovis*, *M. bovis* BCG, and *M. africanum* of the TB complex were differentiated from MOTT based on the different restriction fragments.

Differentiation of fast-growing mycobacteria

20 Lanes 12-16 in Fig. 3 indicated that 644-bp fast-growing mycobacteria including *M. fortuitum* 6841, *M. abscessus*, *M. chelonae*, and *M. peregrinum* were not cleaved by Xho-I, thereby differentiating them from the other mycobacteria (Figs. 3 and 4).

Differentiation of slow-growing bacteria including *M. avium* complex and *M. kansasii*

It was reported that *M. avium* complex and *M. kansasii* were isolated from a clinical sample at the highest frequency in fast-growing bacteria. As shown in Fig. 3, the *M. avium* complex including *M. avium* (lane 5) and *M. intracellulare* (lane 6) produced three kinds of restriction fragments of 391-bp, 169-bp, and 84-bp by treating with the *Xho*I, thereby differentiating them from *M. kansasii* (lane 11) producing two kinds of fragments (391-bp and 253-bp). Therefore, the PRA method according to the present invention can differentiate the reference strains with 100% sensitivity and specificity

Based on the above results, the differentiation of 27 kinds of reference strains including *M. tuberculosis*, *M. bovis*, *M. bovis* BCG, and *M. africanum* of the TB complex, and *M. avium*, *M. intracellulare*, and *M. kansasii* among MOTT are summarized in Fig. 4.

EXAMPLE 6

Differentiation of clinically isolated strains of mycobacteria by using the PRA

The PCR amplification of EXAMPLE 3 and PRA method of EXAMPLE 5 were performed for 198 clinically-isolated strains shown in Table 4.

[Table 4]

PRA analysis of clinically isolated strains

strain	No. of isolates
TB complex	
<i>M. tuberculosis</i>	54
<i>M. bovis</i>	9
Slow-growing mycobacteria	
<i>M. avium</i> complex	49
<i>M. kansasii</i>	30
<i>M. szulgai</i>	12
<i>M. gordonae</i>	9
<i>M. marinum</i>	3
Rapid-growing mycobacteria	
<i>M. fortuitum</i>	17
<i>M. chelonae</i>	15
Sum: 198	

5 Fig. 5 is a photograph of 2% agarose gel electrophoresis of a reaction product obtained by treating the amplified hsp 65 gene fragment with Xho I. In panel A, lane M is a DNA size marker obtained by treating 174 with *Hae* III; Tbc –M of Lanes 1-8 and Mac of Lanes 9-16 are concerned with a result of clinical isolates of *M. tuberculosis* and *M. avium* complex, respectively. In panel B, Kac of lanes 1-8 and Foc of lanes 9-16 represent a result of clinical isolates of *M. kansasii* and *M. fortuitum*, respectively.

10

From Fig. 5, it can be seen that 54 strains of TB complex produced specific restriction fragments of 391bp, 150bp, and 103-bp, thereby differentiating them from 144 strains of MOTT. 32 strains of fast-growing mycobacteria were not cleaved by the restriction enzyme, so they could be differentiated from the other 168 strains. 49 clinical isolates of *M. avium* complex produced specific restriction fragments of 391bp, 169bp, and 84bp, and thus were differentiated from 39 clinical isolates of a group including *M. kansasii* which produced 2 kinds of restriction fragments (391-pb and 253-bp).

10 This example confirmed that the PRA method of the present invention by using the hsp 65 gene can be applied to a clinical isolate of mycobacteria.

EXAMPLE 7

15 Identification of clinical isolate by using the comparative sequence analysis

As shown in Table 2, 38 mycobacterial species including 10 kinds of TB complex and 28 MOTT obtained from the Koran Institute of Tuberculosis (Seoul, Korea) were employed as clinically isolated species.

20 DNA extraction, amplification, and PCR-mediated sequencing of hsp 65 gene fragments were accomplished according to the methods described in Examples 3 and 4. Then, the result was multi-aligned with a sequence database of 54 kinds of reference species in the Megalign program of

Dnastar software according to the method of Example 4, to infer a phylogenetic tree according to the Neighbor-Joining method of Mega software. The identification result showed that 38 clinically isolated strains were identified to species level with 100% sensitivity and specificity as indicated in Table 2 and Figs. 7a to 7d. The result in Figs. 7a to 7d are specifically described:

Fig. 7a: Identification of 4 strains of *M. gordonae* (KIT 32101, 32104, 32105, 32106), 4 strains of *M. szulgai* (KIT 31102, 31103, 31106, 31107), and a strain of *M. marinum* (KIT 21101).

10 Fig. 7b: Identification of 2 strains of *M. scrofulaceum* (KIT 30101, 30102), and 4 strains of *M. avium* complex (KIT 41105, 41110, 41111, 41115).

Fig. 7c: Identification of 1 strain of *M. tuberculosis* (KIT 77710), 3 strains of *M. kansasii* (KIT 20118, 20119, 20120), and 4 strains of *M. terrae* complex (KIT 47101, 47102, 47103, 47104).

Fig. 7d: Identification of 2 strains of *M. chelonae* complex (KIT 61104, 61105), and 4 strains of *M. fortuitum* (KIT 60108, 60109, 60110, 60111).

a) 10 strains of *M. tuberculosis* were identified as *M. tuberculosis* (Table 2 and Fig. 7b). The clinically isolated strains had 100% nucleotide sequence homology with *M. tuberculosis* ATCC 27284.

b) 3 strains of *M. avium* complex were identified as *M. intracellulare* and a strain of *M. avium* complex was identified as *M. avium*, respectively. *M. avium* (KIT 41110) had 99.5% nucleotide sequence homology with *M. avium*

ATCC 25281 which included 3 different nucleotides in a 604-bp hsp 65 gene fragment. When the nucleotide sequences of 3 strains of *M. intracellulare* (KIT 41105, 41111, and 51115) were compared with that of *M. Intracellulare* ATCC 13850, they showed 99.0-99.8% sequence homology.

5 c) 2 strains of *M. scrofulaceum* (KIT 30101, 30102) were identified as *M. scrofulaceum* (Fig.7b). 2 clinically isolated strains and *M. scrofulaceum* ATCC 19981 had 99.8-100% nucleotide sequence homology.

 d) 3 strains of *M. kansasii* (KIT 20118, 20119, 20120) had 100% nucleotide sequence homology, and were identified as *M. kansasii* Type I
10 ATCC 12478 (Fig. 7c). The result is consistent with that of the biochemical identification method.

 e) 4 strains (KIT 32101, 32104, 32105, and 32106) were identified as *M. gordonae* (Fig. 7a and Table 2). When comparing the nucleotide sequences of a 604-bp hsp 65 gene fragment of 4 clinically isolated strains,
15 they had 99.2-99.8% sequence homology with each other, but they had 95.9-96.3% sequence homology with *M. gordonae* ATCC 14470. The result indicates a considerably low sequence homology between *M. gordonae* species.

 Like the results of the biochemical identification method, 4 clinically
20 isolated strains (KIT 31102, 31103, 31106, 31107) were identified as *M. szulgai* (Fig. 7a, and Table 2), and had 99.5-100% sequence homology with their reference strain (*M. szulgai* ATCC 35799)

 Like the results of the biochemical identification method, a clinically

isolated strain was identified as *M. marinum* (Fig. 7a and Table 2), and had 99.3% sequence homology with its reference strain (*M. marinum* ATCC 927). 4 strains were identified as *M. nonchromogenicum* of *M. terrae* complex, and had 95.0-100% sequence homology with *M. nonchromogenicum* ATCC 19530. The result is consistent with the previous report that *M. terrae* complex had sequence heterogeneity.

f) Like the biochemical identification method, 2 clinically isolated strains (KIT 61104, 61105) were identified as *M. abscessus* of *M. chelonae* complex. The identification method of the present invention resolves the problem of the conventional biochemical method in which *M. chelonae* and *M. abscessus* cannot be differentiated. The strains have 98.4-99.5% nucleotide sequence homology with *M. abscessus* CAP97E-03.

Like the biochemical identification method, 4 clinically isolated strains were identified as *M. fortuitum* ATCC 6841, and had 99.4-100% sequence homology with the reference strain. In the present invention, *M. fortuitum* ATCC 6841, *M. fortuitum* ATCC 49403, *M. fortuitum* ATCC 49404, and *M. peregrinum* were employed as reference strains for *M. fortuitum*.

WHAT IS CLAIMED IS:

1. A pair of primers for specifically amplifying an hsp 65 (Heat Shock Protein 65) gene fragment of mycobacterial species comprising the
5 nucleotide sequences as shown in SEQ ID NO: 55 and SEQ ID NO: 56.
2. A polynucleotide of an hsp 65 gene fragment of mycobacterial species wherein the fragment is amplified by using a pair of primers specifically amplifying the hsp 65 gene fragment of mycobacterial species
10 comprising the nucleotide sequences as shown in SEQ ID NO: 55 and SEQ ID NO: 56.
3. A polynucleotide selected from the group consisting of polynucleotides as shown in SEQ ID NO: 1 to SEQ ID NO: 54, and
15 polynucleotides complementary thereto.
4. A polynucleotide set for the detection or identification of mycobacterial species wherein the set comprises at least an hsp 65 gene fragment selected from the group consisting of the polynucleotides as shown
20 in SEQ ID NO: 1 to SEQ ID NO: 54 and polynucleotides complementary thereto.

5. A method for the identification of mycobacterial species comprising the steps of:

(1) amplifying an hsp 65 gene fragment of mycobacterial species of interest with primers for specifically amplifying the hsp 65 gene fragment;

5 (2) analyzing a nucleotide sequence of the amplified hsp 65 gene fragment; and

(3) comparing the nucleotide sequence of the amplified hsp 65 gene fragment obtained in step (2) with a 604-bp hsp 65 gene fragment of a reference mycobacterial species.

10

6. The method of claim 5, wherein the primers comprise the polynucleotides as shown in SEQ ID NO: 55 and SEQ ID NO: 56.

7. The method of claim 5, wherein step (3) of comparing the nucleotide
15 sequence of the mycobacterial species of interest with that of a reference mycobacterial species is performed by multi-aligning the nucleotide sequence of the 604-bp hsp 65 gene fragment of the mycobacterial species of interest with a polynucleotide set of claim 4 to infer a phylogenetic tree.

20 8. A method for the identification of mycobacterial species comprising the steps of:

(1) amplifying an hsp 65 gene fragment of mycobacterial species with primers of claim 1; and

(2) analyzing the amplified fragment according to the RFLP (Restriction Fragment Length Polymorphism) analysis method using a restriction enzyme recognition site in the amplified fragment.

5 9. The method of claim 8, wherein the restriction enzyme is Xho I.

10. The method of claim 9 comprising the step of treating the amplified hsp 65 gene fragment with Xho I to produce restriction fragment(s), and analyzing the restriction fragment(s) according to an RFLP analysis method
10 to differentiate TB complex (*Mycobacterium tuberculosis* complex) and MOTT (*Mycobacteria* other than *Mycobacterium tuberculosis*).

11. The method of claim 10, wherein the restriction fragments are 391-bp, 150-bp, and 103-bp fragments to identify the TB complex.

15

12. The method of claim 10, wherein the 644-bp hsp 65 gene fragment is not cleaved by a restriction enzyme to identify fast-growing mycobacteria of MOTT.

20 13. The method of claim 10, wherein the restriction fragments are 391-bp, 169-bp, and 48-bp to identify a mycobacterial species selected from the group consisting of *M. avium*, *M. intracellulare*, *M. celatum*, *M. shimoidei*, and *M. szulgai*.

14. The method of claim 10, wherein the restriction fragments are 391-bp and 253-bp to identify a mycobacterial species selected from the group consisting of *M. gastri*, *M. genavense*, *M. gordonae*, *M. haemophilum*, *M. kansasii*, *M. malmoense*, *M. marinum*, *M. scrofulaceum*, *M. simiae*, and *M. ulcerans*.
15. A kit for the differentiation or diagnosis of TB complex and MOTT comprising a pair of primers of claim 1 and Xho I, wherein the mycobacterial species is differentiated or diagnosed based on the size of restriction fragment(s) which is obtained by amplifying an hsp 65 gene fragment of mycobacterial species in a sample with the primers to produce an amplified fragment and analyzing the amplified fragment according to an RFLP analysis method.
16. A method for the identification of a mycobacterial species comprising the steps of:
- (1) amplifying an hsp 65 gene fragment of a mycobacterial species of interest with primers for specifically amplifying an hsp65 gene of mycobacteria; and
 - (2) hybridizing the amplified hsp65 gene fragment with a probe set comprising at least a probe selected from the group consisting of polynucleotides as shown in SEQ ID NO: 1 to SEQ ID NO: 54 and polynucleotides complementary thereto.

FIG.1

1623-bp *hsp65* gene






FIG.2




3/8

FIG.3




BEST AVAILABLE COPY


FIG.4



5/8

FIG.5



6/8
FIG.6Slow-growing
mycobacteriaFast-growing
mycobacteria

Outgroup

Scale: each - is approximately equal to the distance of 1.366123



7/8
FIG.7A

FIG.7B





8/8
FIG.7C

FIG.7D



Sequence Listing

<110> KIM, Bum-Joon
BIOMEDLAB CORPORATION

<120> PRIMERS FOR AMPLIFYING HSP 65 GENE OF MYCOBACTERIAL SPECIES, HSP
65 GENE FRAGMENTS AND METHOD OF IDENTIFYING MYCOBACTERIAL SPECIES
WITH THE SAME

<130> OPP021096KR

<150> KR 10-2002-0004297
<151> 2002-01-24

<150> KR 10-2002-0011648
<151> 2002-03-05

<160> 56

<170> KopatentIn 1.71

<210> 1
<211> 604
<212> DNA
<213> Mycobacterium abscessus

<400> 1

ggaggaccgc tacgagaaga tcggcgctga gctgggtcaag gaagtgcca agaagaccga	60
cgacgtcgcg ggtgacggca ccaccaccgc caccgtgctc gccaggctc tggtaagga	120
aggtctgcgt aacgtcgccg ccggcgccaa cccgctcggc ctgaagcgcg gtatcgagaa	180
ggccgtcgag aaggtcaccg agacgtgct gaagagcgc aaggagtcg agaccaagga	240
gcagatcgcg gccacggccg gtatctccgc gggcgaccag tccatcggcg acctgatcgc	300
cgaggccatg gacaaggttg gtaacgaggg tgatcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctggagctca ccgaggggat gcgcttcgac aagggtaca tctcgggcta	420
cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gatccctaca tcctgctggt	480

Sequence Listing

cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggtctctt ccactctggt 600

cgtc 604

<210> 2

<211> 604

<212> DNA

<213> *Mycobacterium africanum*

<400> 2

ggaggatccg tacgagaaga tcggcgccga gctgggtcaaa gaggtagcca agaagaccga 60

tgacgtcgcc ggtgacggca ccacgacggc caccgtgctg gccagggcgt tggttcgca 120

gggcctgcgc aacgtcgcg ccggcgccaa cccgctcggc ctcaaacgcg gcatcgaaaa 180

ggcgtggag aaggtcaccg agaccctgct caagggcgcc aaggaggctg agaccaagga 240

gcagattgcg gccaccgcag cgatttcggc ggggtgaccag tccatcggtg acctgatcgc 300

cgaggcgatg gacaagggtg gcaacgaggc cgtcatcacc gtcgaggagt ccaacacctt 360

tgggctgcag ctcgagctca ccgaggggat gcggttcgac aagggtaca tctcggggta 420

cttcgtgacc gaccggagc gtcaggagc ggtcctggag gaccctaca tcctgctggt 480

cagctccaag gtgtccactg tcaaggatct gctgccgctg ctcgagaagg tcacggagc 540

cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgtgt ccaccctggt 600

cgtc 604

<210> 3

<211> 604

<212> DNA

<213> *Mycobacterium asiaticum*

Sequence Listing

<400> 3

ggaggacccg tacgagaaga tcggcgccga gctgggtcaag gaagtcgcca agaagaccga	60
cgacgtggcc ggtgacggca ccacgacggc caccgtgctg gcacaggcgc tggtaagga	120
gggcctgcgc aacgttgccg caggcgccaa cccgctgggc ctgaagcgcg gcatcgagaa	180
ggccgtcgag aaggtcaccc agaccctgct cagctcggcc aaggacgtcg agaccaagga	240
gcagatcgcg gccaccgcgg gtatttcgcg gggcgaccag tcgatcggcg acctgatcgc	300
cgaggcgatg gacaaagtcg gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgaggcat gcggttcgac aagggttaca tctcgggcta	420
cttcgtcacc gacgccgagc gtcaggaagc cgtcctggag gaccctaca tcctgctggt	480
ttccagcaag gtgtcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccaggc	540
cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt	600
cgtc	604

<210> 4
 <211> 604
 <212> DNA
 <213> *Mycobacterium aichiense*

<400> 4

cgaggacccg tacgagaaga tcggcgctga gctgggtcaag gaagtcgcca agaagactga	60
cgatgtcgcg ggcgacggca ccaccaccgc caccgtgctc gtcaggctc tggttcgca	120
aggtctgcgc aacgtcgtg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggccgtcgag aagatcaccc agacgtcct caagagcgc aaggaggtcg agaccaagga	240
ccagatcgcg gccaccgcgg gcatctcgc gggcgaccag accatcggtg acctgatcgc	300

Sequence Listing

cgaggccatg gacaagggtcg gcaacgaggg tgtcatcacc gtcgaggagt cgaacacctt	360
cggcctgcag ctcgagctca ccgaggggtat gcgcttcgac aagggctaca tctcgggtta	420
cttcgtgacc gacgccgagc gtcaggaagc ggtcctcgag gatccgtaca tcctgctggt	480
gtcgtcgaag gtctcgaccg tcaaggacct gcttccttg ctggagaagg tcattcagtc	540
gggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctgt ccaccctggt	600
ggtc	604
<div style="display: flex; justify-content: space-between;"> <div style="width: 40%;"> <p><210> 5</p> <p><211> 604</p> <p><212> DNA</p> <p><213> Mycobacterium avium</p> </div> <div style="width: 55%;"></div> </div>	
<div style="display: flex; justify-content: space-between;"> <div style="width: 40%;"> <p><400> 5</p> </div> <div style="width: 55%;"> <p>ggaggaccg tacgagaaga tcggcgccga gctgggtcaag gaagtcgcca agaagaccga</p> <p>cgacgtcgcc ggtgacggca cgacgacggc cacgggtgtc gcccaggcgt tgggccgga</p> <p>gggcctgcgc aacgtcgcg ccggcgccaa cccgctgggt ctcaagcgcg gcatcgagaa</p> <p>ggccgtcgag aagggtaccg agaccctgct caagtcggcc aaggaggtcg agaccaagga</p> <p>ccagatcgct gccaccgcgg ccattctcgc gggcgaccag tcgatcggcg acctgatcgc</p> <p>cgaggcgatg gacaagggtcg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt</p> <p>cggcctgcag ctcgagctca ccgaggggtat gcggttcgac aagggttaca tctcgggcta</p> <p>cttcgtcacc gacgccgagc gtcaggaagc cgtcctcgag gatccgttca tcctgctggt</p> <p>cagctccaag gtctcgaccg tcaaggacct gctgccgctg ctggagaagg tcattccaggc</p> <p>cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggccctgt ccaccctggt</p> <p>cgtc</p> </div> </div>	<div style="display: flex; justify-content: space-between;"> <div style="width: 40%;"></div> <div style="width: 55%;"> <p>60</p> <p>120</p> <p>180</p> <p>240</p> <p>300</p> <p>360</p> <p>420</p> <p>480</p> <p>540</p> <p>600</p> <p>604</p> </div> </div>

Sequence Listing

<210> 6
 <211> 604
 <212> DNA
 <213> Mycobacterium bovis

<400> 6
 ggaggatccg tacgagaaga tcggcgccga gctgggtcaaa gaggtagcca agaagaccga 60
 tgacgtcgcc ggtgacggca ccacgacggc caccgtgctg gcccaggcgt tggttcgca 120
 gggcctgcgc aacgtcgcg cggcgccaa cccgctcggc ctcaaacgcg gcatcgaaaa 180
 ggccgtggag aaggtcaccg agaccctgct caagggcgcc aaggaggctc agaccaagga 240
 gcagattgcg gccaccgag cgatttcggc gggtgaccag tccatcggtg acctgatcgc 300
 cgaggcgatg gacaaggtgg gcaacgaggc cgtcatcacc gtcgaggagt ccaacacctt 360
 tgggctgcag ctcgagctca ccgagggtat gcggttcgac aagggctaca tctcggggta 420
 cttcgtgacc gacccgagc gtcaggaggc ggtcctggag gaccctaca tcctgctggt 480
 cagctccaag gtgtccactg tcaaggatct gctgccgctg ctcgagaagg tcatcgagc 540
 cggtaaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt 600
 cgtc 604

<210> 7
 <211> 604
 <212> DNA
 <213> Mycobacterium bovis BCG

<400> 7
 ggaggatccg tacgagaaga tcggcgccga gctgggtcaaa gaggtagcca agaagaccga 60
 tgacgtcgcc ggtgacggca ccacgacggc caccgtgctg gcccaggcgt tggttcgca 120

Sequence Listing

gggcctgcgc aacgtcgcg ccggcgccaa cccgctcggc ctcaaacgcg gcatcgaaaa	180
ggccgtggag aaggtcaccg agaccctgct caagggcgcc aaggaggtcg agaccaagga	240
gcagattgcg gccaccgcag cgatttcggc gggtgaccag tccatcggcg acctgatcgc	300
cgaggcgatg gacaaggtgg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
tgggctgcag ctcgagctca ccgaggggat gcggttcgac aagggtctaca tctcggggta	420
cttcgtgacc gaccgggagc gtcaggaggc ggtcctggag gacccttaca tctgctggt	480
cagctccaag gtgtccactg tcaaggatct gctgccgctg ctcgagaagg tcatcgagc	540
cggtaaagcg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt	600
cgtc	604

<210> 8
 <211> 604
 <212> DNA
 <213> Mycobacterim celatum Type 1

<400> 8 ggaggacccc tacgaaaaga tcggcgccga gctggtaag gaagtcgcca agaagaccga	60
cgacgtcgcg ggtgacggta cgacgacggc cacggtgctg gccagggcg tggtaagga	120
gggcctgcgc aacgtcgccg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggccgtcgag aaggtcaccg agacgtgct caagggcgcc aaggaggtcg agaccaagga	240
gcagattgct gccaccgcg ccatctccgc cggcgaccag tcgatcgcg acctgatcgc	300
cgaggccatg gacaaggtcg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgaggggat gcggttcgac aagggtctaca tctcgggtta	420

Sequence Listing

cttcgtcacc gacgccgagc gtcaggaggc ggtgctcgag gagccgtaca tcctgctggt	480
cagctccaag gtgtcgacgg tcaaggacct gcttccgctg ctggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctct ccaccctggt	600
cgtc	604

<210> 9
 <211> 604
 <212> DNA
 <213> Mycobaterium celatum TypeII

<400> 9	
ggaggacccc tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga	60
cgacgtcgcg ggtgacggta cgacgacggc caccgtgctg gcccaggcgc tggtaagga	120
aggcctgcgc aacgtcgccg ccggtgccaa cccgctcggc ctgaagcgcg gtatcgagaa	180
ggccgtcgag aaggtcaccg agacgtgct caagggcgcc aaggaggtcg agaccaagga	240
gcagatcgct gccaccgcgg ccatctccgc cggtgaccag tcgatcggcg acctgatcgc	300
cgaggcgatg gacaaggctg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgagggat gcgcttcgac aagggtaca tctcgggtta	420
cttcgtcacc gacgccgagc gtcaggaggc ggtgctcgag gagccctaca tcctgctggt	480
cagctccaag gtgtcgacgg tcaaggatct gctgccgctg ctggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgttga gcaccctggt	600
cgtc	604

<210> 10
 <211> 604

Sequence Listing

<212> DNA

<213> *Mycobacterium chelonae*

<400> 10

ggaggacccg tacgagaaga tcggcgctga gctggtcaag gaagttgcca agaagactga	60
cgacgtcgcg ggtgacggca ctactaccgc caccgtgctt gcccaggctc tggtaagga	120
aggtctgcgt aacgtcgctg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggccgtggag gccgtcacca gctctctgct ggactccgcc aaggagatcg acaccaagga	240
gcagatcgcg gccaccgcgg gcatctccgc gggtgaccag tccatcggtg atctgatcgc	300
cgaggccatg gacaaggctg gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctggagctca ccgagggcat gcgcttcgac aagggtaca tctcgggtta	420
cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gatccctaca tcctgctggt	480
cagctccaag gtctcgaccg tcaaggacct acttccttg ctggagaagg tcatccaggg	540
cggaagccg ctgctgatca tcgccgagga cgttgagggc gaggtctctt cgaccctggt	600
cgtc	604

<210> 11

<211> 604

<212> DNA

<213> *Mycobacterium chitae*

<400> 11

ggaggacccg tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagactga	60
cgacgtcgcc ggcgacggca ccaccaccgc caccgttctg gcccaggcgc tggttcgca	120
aggtctgcgc aacgtcgcg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggccgtcgag accgtctcgg agaacctgct caagtcggcc aaggaggtcg agaccaagga	240

Sequence Listing

gcagatcgcc gccaccgccg ggatctccgc gggcgacacc accatcgggtg acctgatcgc	300
cgaggccatg gacaaggtgg gcaacgaggg tgatcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctggagctca ccgagggcat gcgcttcgac aagggtctaca tctcgggcta	420
cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gatccctaca tcctgctggt	480
cagctcgaag atctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc	540
cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcctgt cgaccctggt	600
ggtc	604

<210> 12
 <211> 604
 <212> DNA
 <213> Mycobacterium microti

<400> 12 ggaggatccg tacgagaaga tcggcgccga gctggtaaaa gaggtagcca agaagaccga	60
tgacgtcgcc ggtgacggca ccacgacggc caccgtgctg gccagggcgt tggttcgcga	120
gggcctgcgc aacgtcgcgg ccggcgccaa cccgctcggg ctcaaacgcg gcatcgaaaa	180
ggccgtggag aaggtcaccg agaccctgct caagggcgcc aaggaggtcg agaccaagga	240
gcagattgcg gccaccgcag cgatttcggc gggtgaccag tccatcgggtg acctgatcgc	300
cgaggcgatg gacaaggtgg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
tgggctgcag ctcgagctca ccgaggggat gcggttcgac aagggtctaca tctcggggta	420
cttcgtgacc gacccggagc gtcaggaggc ggtcctggag gaccctaca tcctgctggt	480
cagctccaag gtgtccactg tcaaggatct gctgccgctg ctcgagaagg tcatcgagc	540

Sequence Listing

cggtaaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt 600

cgtc. 604

<210> 13

<211> 604

<212> DNA

<213> *Mycobacterium flavescens*

<400> 13

ggaggaccgc tacgagaaga tcggcgctga gctggtaag gaagtcgcca agaagaccga 60

cgacgtcgcg ggcgacggca ccaccaccgc caccgtgctg gcccaggcgc tcgtgcgcga 120

gggtctgcgc aacgtcgcg cggcgccaa cccgatggcg ctgaagcgcg gtatcgagaa 180

ggccgtcgag aaggtcaccg agacgtgct gaagtcggcc aaggaggtcg agaccaagga 240

gcagatcgct gccaccgcgc cgatctcggc gggcgacacc cagatcgga agctgatcgc 300

cgaggccatg gacaaggctg gcaacgaggg tgtcatcacc gttgaggagt ccaacacctt 360

cgggctgcag ctcgagctca ccgagggat gcgcttcgac aagggtaca tctcgggtta 420

cttcgtgacc gacgccgagc gtcaggaagc ggtcctcgag gatccctgca tcctgctcgt 480

gtcgtccaag gtgtcgaccg tcaaggatct gtcgccgttg ctggagaagg tcattcagga 540

cggcaagccg gtgctgatca tcgccgagga cgtcgagggg gaggccctgt cgaccctggt 600

ggtc 604

<210> 14

<211> 604

<212> DNA

<213> *Mycobacterium fortuitum* 6841

<400> 14

Sequence Listing

ggaggacccg tacgagaaga tcggcgctga gctcgtcaaa gaggtcgcca agaagaccga	60
cgacgtcgcg ggcgacggca ccaccaccgc caccgttctg gcacaggccc tggttcgtga	120
aggtctgcgc aacgtcgctg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggccgtcgag aaggtcaccg agacgtgct gaagagcgcc aaggaggtgg agaccaagga	240
gcagatcgct gccaccgccg gtatctccgc cggtgaccag tccatcggtg acctgatcgc	300
cgaggccatg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggaga gcaacacctt	360
cggcctgcag ctggagctca ccgggggtat gcgcttcgac aagggtaca tctcgggcta	420
cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gatccctaca tcctgctggt	480
cagctccaag gtctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc	540
cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctgt cgaccctggc	600
ggtc	604

<210> 15
 <211> 604
 <212> DNA
 <213> Mycobacterium fortuitum 49403

ggaggacccg tacgagaaga tcggcgctga gctcgtcaaa gaggtcgcca agaagaccga	60
cgacgtcgcg ggcgacggca ccaccaccgc caccgttctg gccaggccc tggttcgca	120
aggtctgcgc aacgtcgctg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggccgtcgag aaggtcaccg agacgtgct gaagagcgcc aaggaggtgg agaccaagga	240
gcagatcgct gccaccgccg gtatctccgc cggtgaccag tccatcggtg acctgatcgc	300
cgaggccatg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggaga gcaacacctt	360

Sequence Listing

cggcctgcag ctggagctca ccgaggggtat gcgcttcgac aagggtctaca tctcgggtta	420
cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gatccctaca tcctgctggt	480
cagctccaag gtctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctgt ccaccctggt	600
ggtc	604
<210> 16	
<211> 604	
<212> DNA	
<213> Mycobacterium fortuitum 49404	
<400> 16	
ggaggaccg tacgagaaga tcggcgaga gctggccaag gaagtcgcca agaagactga	60
cgacgtcgca ggcgacggca ccaccacggc caccgtgctc gccaggctc tggttcgca	120
aggtctgcgc aacgtcgag ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggctgtcggg gccgtcacc agacgtgct gaagtcgcc aaggaggagg agaccaagga	240
gcagatcgt gccaccgcc cgatctcgc cggtgacgtc cagatcggcg agctcatcgc	300
cgaggccatg gacaaggctg gcaacgagg tgatcatcacc gtcgaggagt cgaacacctt	360
cggcctgcag ctggagctca ccgaggggtat gcgcttcgac aagggtctaca tctcgggtta	420
cttcgtgacc gacgccgagc gtcaggaagc ggtcctcgag gatccgtaca tcctgctcgt	480
ctcgtcgaag gtctcgacgg tcaaggacct gctgccctg ctggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctgt ccaccctggt	600
ggtc	604

Sequence Listing

<210> 17
 <211> 604
 <212> DNA
 <213> *Mycobacterium gastr*

<400> 17
 ggaggacccg tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga 60
 cgacgtcgcc ggcgacggca ccaccacggc caccgtgctc gcgcaggcgc tggtaagga 120
 gggcctgcgc aacgtcgcg cggcgccaa cccgctgggc ctgaagcgcg gcatcgagaa 180
 ggccgtcgag aaggtcaccg agacgtgct caagggcgcc aaggaggtcg agaccaagga 240
 gcagatcgcg gccaccgcgg ccatctccgc cggtgaccag tcgatcgcg acctgatcgc 300
 cgaggcgatg gacaaggtgg gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt 360
 cggcctgcag ctcgagctca ccgagggcat gcggttcgac aagggtaca tctccggcta 420
 ctctgtcacc gacgtgagc gtcaggaagc tgttctggag gaccctaca tcctgctggt 480
 cagctcgaag gtctcgaccg tcaaggacct gctgccgctg ttggagaagg tcatccaggc 540
 gggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt 600
 cgtc 604

<210> 18
 <211> 604
 <212> DNA
 <213> *Mycobacterium genavense*

<400> 18
 ggaggacccc tacgagaaga tcggcgctga gctggtcaag gaagtcgcca agaagaccga 60
 cgacgtcgcc ggtgacggca ccacgacggc caccgtgctc gtcaggcgc tcgtcaagga 120

Sequence Listing

gggcctgcgc aacgtggcgg ccggcgccaa cccgctgggc ctcaagcgcg gcatcgagaa	180
ggccgtcgaa aaggtcaccg agacgtgct gaagtcggcc aaggatgtcg agaccaagga	240
ccagatcgct gccaccgccg cgatttcgcg gggcgaccag tcgacggcg acctgatcgc	300
cgaggcgatg gacaaggctg gcaacgaggc cgtcatcacc gtcgaggagt ccaacacctt	360
cgggctgcag ctcgagctca ccgaggggat gcgcttcgac aagggtaca tctcgggcta	420
cttcgtcacc gacgccgagc gtcaggaagc cgtcctggag gacccttca tcctgctggt	480
cagctccaag gtgtcgacgg tcaaggacct gctgccgtg ctggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgtga gcaccctggt	600
cgtc	604

<210> 19
 <211> 603
 <212> DNA
 <213> Mycobacterium gordonae

<400> 19 gaggacccgt acgagaagat cggcgctgag ctggtcaagg aagtcgcca gaagaccgac	60
gacgttgccg gcgacggcac gacgacggcg accgtgctgg cgcaggcact ggtcaaggaa	120
ggcctgcgca acgtagccgc cggcgccaac ccgctggggc tgaagcgcg catcgagaag	180
gccgtggaga aggtcaccca gaccctgctc agctcggcca aggacgtcga gaccaaggag	240
cagatcgcg ccaccgcggg catctccgcg ggtgaccagt cgatcgggta cctgatcgcc	300
gaggcgatg acaaggctcg caacgagggc gtcacaccg tcgaggagtc caacaccttc	360
ggcctgcagc tcgagctgac cgagggcatg cggttcgaca agggctacat ctcgggctac	420
ttcgtcaccg acgccgagc tcaggaagcc gtcctggaag acccctacat cctgctggtg	480

Sequence Listing

tccagcaagg tgctgaccgt gaaggacctg ctgccgctgc tggagaaggt cattcagggt	540
ggcaagccgc tgctgatcat cgccgaggac gtcgagggcg aagcgctgtc gaccctggtc	600
gtc	603

<210> 20
 <211> 604
 <212> DNA
 <213> Mycobacterium haemophilum

<400> 20 ggaggacccg tacgagaaga tcggcgccga gctgggtcaag gaagtcgccga agaagaccga	60
cgacgtcgct ggtgatggca ccacgacggc gacggtgctg gtcaggcgcg tggtaaaga	120
gggcctgcgt aacgtcgcg cggcgccaa cccgctgggt ctcaagcgcg gcatcgagaa	180
ggcggtcgag aagatcaccg agacgtgct caagggcgcc aaggaggctg agaccaagga	240
ccaaattgcg gccaccgag cgatctcggc gggtgaccag tcgatcggcg acctgatcgc	300
cgaggcgatg gacaaggctg gcaacgagg cgatcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgagggcat gcggttcgat aagggtaca tctcgggcta	420
cttcgtcacc gacgccgagc gccaggaagc cgtcctggag gaccctaca tcctgctggt	480
cagctccaag gtgtcgaccg tcaaggacct gctgccactg ttggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt	600
cgtc	604

<210> 21
 <211> 603
 <212> DNA

Sequence Listing

<213> Mycobacterium interjectum

<400> 21

gaggacccgt acgagaagat cggcgccgag ctggtcaagg aagtcgcaa gaagaccgac	60
gacgtcgccg gtgacggcac gacgacggcc acggtgctgg cccaggcctt ggtcaaggag	120
ggcctgcgca acgtcgcggc cggcgccaac ccgcccggcg tcaagcgcg catcgaaaag	180
gccgtcgaga aggtcaccga gaccctgctg aagtcggcca aggatgtcga gaccaaggag	240
cagatcgccg cgaccggcgc gatctccgcg ggcgaccagt cgatcgcgga cctcatcgcc	300
gaggcgatgg acaaggctcg caacgagggc gtcatcaccg tcgaggagtc caacaccttc	360
ggcctgcagc tcgagctcac cgagggcatg cggttcgaca agggctacat ctcggggtac	420
ttcgtcaccg acgccgagcg tcaggaagcg gtccctcgagg acccctacat cctgctggtc	480
agctcgaagg tgtcgacggt caaggacctg ttgccgctgc tggagaagg catccaggcc	540
ggcgagccgc tgttgatcat cgccgaggac gtcgagggcg aggcgctgtc caccctggtc	600
gtc	603

<210> 22

<211> 604

<212> DNA

<213> Mycobacterium intermedium

<400> 22

ggaggacccg tacgagaaga tcggcgccga gctggtcaag gaagttgcca agaagacgga	60
cgacgtcgcc ggtgacggca ccacgacggc caccgtgtc gccagggcg tggcgcgga	120
gggtctgcgc aatgtcgctg ccggtgcaa cccgctgagc ctgaagcgcg gtatcgagaa	180
ggcagtcgag aaggtcaccg agaccctgct caagtcggcc aaggaggctg agaccaagga	240

Sequence Listing

ccagatcgct gccaccgcag cgatttccgc gggggaccag tcgatcggcg acctgatcgc	300
cgaggcgatg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag cttgagctca ccgaggggat gcggttcgac aagggttaca tctcgggcta	420
cttcgtcacc gacgccgagc gtcaggaagc cgtcctggaa gaccctgata tctgctggt	480
cagctccaag gtttcgacgg tcaaggacct gctcccgtg ctggagaagg tcattcaggc	540
cggaagccg ctgctgatca tcgccgagga cgtcagggc gaggcgtga gcaccctggt	600
cgtc	604

<210> 23
 <211> 604
 <212> DNA
 <213> Mycobacterium intracellulare

<400> 23 ggaggaccg tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga	60
cgacgtcgcc ggtgacggca cgacgacggc cacggtgctg gtcagggcgt tggcccgca	120
gggcctgcgt aacgtcgccg ccggcgccaa cccgctgggt ctcaagcgcg gcatcgagaa	180
ggccgtcgag aaggtcaccg agaccctgct caagtcggcc aaggaggtcg agaccaagga	240
ccagatcgct gccaccgcg cgatttcggc gggcgaccag tcgatcggcg acctcatcgc	300
cgaggggatg gacaaggtcg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgagggcat gcggttcgac aagggttaca tctcgggcta	420
cttcgtcacc gacgccgagc gtcaggaagc ggtcctcgag gacccttca tctgctggt	480
cagctccaag gtgtcgacgg tcaaggacct gctgccgtg ctggagaagg tcattcaggc	540
cggaagccg ctgctgatca tcgccgagga cgtcagggc gaggcctga gcaccctggt	600

Sequence Listing

	cgtc	604
<210>	24	
<211>	604	
<212>	DNA	
<213>	Mycobaterium kansasii Type I	
<400>	24	
	ggaggacccg tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga	60
	cgacgtcgct ggcgacggca ccaccacggc caccgtgctt gcgcaggcgc tggtaaaga	120
	gggcctgcgc aacgtcgcg ccggcgccaa cccgctgggc ctcaagcgcg gcatcgagaa	180
	ggccgtcgag aaggtcaccg agacgtgct caagggcgcc aaggaggtcg agaccaagga	240
	gcagatcgcg gcgaccggc ccatctccgc cggcgaccag tcgatcggcg acctgatcg	300
	cgaggcgatg gacaagggtc gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt	360
	cggcctgcaa ctcgagctca ccgagggcat gcggttcgac aagggttaca tctccggcta	420
	cttcgtcacc gacgccgagc gtcaggaagc ggttctggag gaccctaca tcctgctggt	480
	cagctcgaag gtatcgacgg tcaaggacct gctgccgctg ctggagaagg tcatccaggc	540
	cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt	600
	cgtc	604
<210>	25	
<211>	604	
<212>	DNA	
<213>	Mycobaterium kansasii Type II	
<400>	25	
	ggaggacccg tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga	60

Sequence Listing

cgacgtcgcc ggcgacggca ccaccacggc cactgtgctc gcgcaggcgt tggtaaaga	120
gggcctgcgc aacgtgcgcg ccggcgccaa cccactgggc ctgaagcgcg gcatcgagaa	180
ggcagtcgag aaggtcaccg agacgctgct caagggcgcc aaggaggtcg agaccaagga	240
gcagatcgct gccaccgcg ccattctcgc gggtgaccag tcgacggcg acctgatcgc	300
cgaggcgatg gacaaggtgg gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgaggtat gcggttcgac aagggtaca tctccggcta	420
cttcgtcacc gacgccgagc gtcaggaagc agttctggag gaccctaca tctgctggt	480
cagctccaag gtgtccaccg tcaaggacct gctgccgctg ctggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt	600
cgtc	604

<210> 26
 <211> 604
 <212> DNA
 <213> Mycobacterium kansasii Type III

ggaggaccg tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga	60
cgacgtcgcc ggcgacggca ccaccacggc cactgtgctc gcgcaggcgc tcgtcaagga	120
gggcctgcgc aacgtggcgg ccggcgccaa cccgctgggc ctgaagcgcg gcatcgagaa	180
ggccgtcgag aaggtcaccg agacctgtt caagggcgcc aaggaggtcg agaccaagga	240
gcagatcgcg gccaccgcg ccattctcgc cggtgaccag tcgattggcg acctgatcgc	300
cgaggcgatg gacaaggtag gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt	360

Sequence Listing

aggcctgcag ctcgagctca ccgagggat gcgctttgac aagggtaca tctccggcta	420
cttcgtcacc gacgccgagc gtcaggaagc agtgctggaa gaccctaca tctgctggt	480
cagctccaag gtgtcgacgg tcaaggacct gctgccgctg ctggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggt gaggctttga gcaccctggt	600
cgtg	604
<div style="display: flex; justify-content: space-between;"> <210> 27 <211> 604 </div> <div style="display: flex; justify-content: space-between;"> <212> DNA <213> Mycobacterium leprae </div>	
<div style="display: flex; justify-content: space-between;"> <400> 27 </div>	
ggaggacccg tacgagaaga ttggcgctga gttggtcaag gaagtcgcca agaagacaga	60
tgacgtcgcc ggtgatggca ccacgacggc caccgtgctg gccaggcat tggtaaaga	120
gggcctacgc aacgtcgcg ccggcgccaa cccgctaggt ctcaagcgtg gcatcgagaa	180
agctgtcgat aaggtaaactg agactctgct caaggacgct aaggaggtcg aaaccaagga	240
acaaattgct gccactgcag cgatttcggc gggtgaccag tcgatcggtg atctgatcgc	300
cgaggcgatg gacaagggtg gcaacgaggg tgttatcacc gtcgaggaat ccaacacctt	360
cggctctgcag ctcgagctca ccgagggat gcggttcgac aagggtaca tttcgggcta	420
cttcgtcacc gacgccgagc gtcaggaagc tgtcctagag gagccctaca tccttctggt	480
cagctccaaa gtgtctaccg tcaaggacct gctgccgctg ctagagaagg tcatccaggc	540
cggcaagtcg ctgctgatca ttgctgagga tgtcgagggt gaggcgttgt ctaccctggt	600
cgtc	604

Sequence Listing

<210> 28
 <211> 604
 <212> DNA
 <213> *Mycobacterium malmoense*

<400> 28
 ggaggacccg tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga 60
 cgacgtggcc ggtgacggca cgacgacggc caccgtgctg ggcagggcg tggtcaaaga 120
 gggcctgcgc aacgtcgcg ccggtgccaa cccgctcagc ctcaagcgcg gcatcgagaa 180
 ggcggtcgag aaggtcaccg agaccctgct caagtcggcc aaggaggtcg agaccaagga 240
 gcagatcgcc gcgaccgccc cgatctcggc gggcgaccag tcgatcggtg acctgatcgc 300
 cgaggcgatg gacaaggctg gcaacgaggg cgtcctcacc gtcgaggagt ccaacacctt 360
 cggcctgcag ctcgagctca ccgagggcat gcggttcgac aagggtaca tctcgggcta 420
 ctctgctacc gaccccgagc gtcaggaagc ggtcctggag gaccctaca tcctgctggt 480
 cagctccaag gtgtcgacgg tcaaggacct gctgccgctg ctggagaagg tcattcaggc 540
 cggcaagccc ctgctgatca tcgccgagga cgtcgagggc gaggcgctct ccaccctggt 600
 cgtc 604

<210> 29
 <211> 604
 <212> DNA
 <213> *Mycobacterium marinum*

<400> 29
 ggaggacccg tacgagaaga tcggcgctga gctggtcaag gaagttgcca agaagaccga 60
 cgacgtggcc ggtgacggca cgacgacggc caccgtgctg gccagggcg tggtcaagga 120
 aggcctgcgc aacgttgcg ccggtgccaa cccgctcggc ctgaagcgcg gcatcgagaa 180

Sequence Listing

ggcagtcgag aaggtcaccg agaccttgct caagtcggcc aaagaggtcg agaccaagga	240
gcagatcgcg gcgaccgag ccatctccgc cggcgaccag tcgatcggcg acctgatcgc	300
cgaggcgatg gacaaggtgg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgaggggat gcggttcgac aagggctaca tctcgggcta	420
cttcgtcacc gacgccgagc gtcaggaagc ggtcctggag gacccctaca tcctgctggt	480
cagttccaag gtgtccaccg tgaaggacct gctgccgctg ctggagaagg tcattcaggg	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt	600
cgtc	604

<210> 30
 <211> 604
 <212> DNA
 <213> Mycobacterium mucogenicum

<400> 30	
ggaggaccg tacgagaaga tcggcgctga gctggtcaag gaagttgcca agaagacgga	60
cgacgtcgtt ggcgacggca ccaccaccgc caccgtgctg gccaggccc tggttcgga	120
aggcctgcgc aacgtcgtg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggcgtcgag gctgtcacca agggcctgct ggcttcgcc aaggaggtcg agaccaagga	240
gcagatcgtt gccaccgccg ggatctcggc cggtgaccag tccatcggcg acctgatcgc	300
cgaggccatg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggaga gcaacacctt	360
cggcctgcag ctggagctca ccgagggat gcgcttcgac aagggctaca tctcgggta	420
cttcgtgacc gacgccgagc gtcaggaagc ggtcctcgag gacccgttca tcctgctggt	480

Sequence Listing

cagctcgaag atctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc 540

gggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctgt cgaccctggt 600

cgtc 604

<210> 31
 <211> 604
 <212> DNA
 <213> Mycobacterium neoaurum

<400> 31

ggaggaccg tacgagaaga tcggcgccga gctgggtcaaa gaggtcgcca agaagaccga 60

tgacgtcgcg ggcgacggca ccaccaccgc caccgtgctg gcccaggccc tggttcgcga 120

aggtctgcgc aacgtcgcg cggcgccaa cccctcggc ctgaagcgcg gcatcgagaa 180

ggcgtcgcg gccgtcaccg agcgcctgct ctcgaccgcc aaagaggtcg agaccaagga 240

gcagatcgct gccaccgcg gcatctccgc cggtgaccag tcgatcggtg acctgatcgc 300

cgaggcgtg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt 360

cggcctgcag ctggagctca ccgagggtat gcgcttcgac aagggtctaca tctcgggtta 420

cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gatccctaca tcctgctggt 480

cagctccaag gtctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc 540

cggcaagccg ttgctgatca tcgccgagga cgtcgagggc gaagccctgt cgaccctggt 600

ggtc 604

<210> 32
 <211> 604
 <212> DNA
 <213> Mycobacterium nonchromogenicum

Sequence Listing

<400> 32
 ggaggatccc tacgagaaga tcggcgctga gctgggtcaaa gaggtcgcca agaagactga 60
 cgacgtcgcg ggtgacggca ccaccaccgc caccgtgctc gcccaggccc tgggtcaagga 120
 aggctcgcg aacgtggccg ccggcgccaa cccgctgggt ctgaagcgcg gcatcgagaa 180
 ggccgttgag aaggtcacct cgaccctgct ggcttcggcc aaggaggtcg agaccaagga 240
 gcagatcgcg gccaccgccg gtatctccgc gggtgaccag agcatcggtg acctgatcg 300
 cgaggccatg gacaaggctg gcaacgaagg tgtcatcacc gtcgaggagt ccaacacctt 360
 cggcctgcag ctggagctca ccgagggcat gcgcttcgac aagggttaca tctcgggtta 420
 cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gacccttaca tcctgctggt 480
 cagctcgaag atctcgaccg tcaaggacct gctgcccttg ctggagaagg tcatccagtc 540
 cggcaagccg ttgctgatca tcgccgagga cgtcgagggc gaggcctgt cgaccctggt 600
 cgtg 604

<210> 33
 <211> 604
 <212> DNA
 <213> Mycobacterium paratuberculosis

<400> 33
 ggaggaccg tacgagaaga tcggcgccga gctgggtcaag gaagtcgcca agaagaccga 60
 cgacgtcgcc ggtgacggca cgacgacggc cacggtgctc gcccaggcgt tggtcgcgca 120
 gggcctcgcg aacgtcgcg ccggcgccaa cccgctgggt ctcaagcgcg gcatcgagaa 180
 ggccgtcgag aaggtcacct agaccctgct caagtcggcc aaggaggtcg agaccaagga 240
 ccagatcgct gccaccgccg ccattctccgc gggcgaccag tcgatcgcg acctgatcg 300

Sequence Listing

cgaggc gatg gacaagg tgc gcaacgagg cgatcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgagggtat gcggttcgac aagggttaca tctcgggcta	420
cttcgtcacg gacgccgagc gtcaggaagc ggtcctcgag gaccggttca tcctgctggt	480
cagctccaag gtctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccaggc	540
cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggccctgt ccaccctggt	600
cgtc	604

<210> 34
 <211> 604
 <212> DNA
 <213> Mycobacterium phlei

<400> 34 cgaggatccg tacgagaaga tcggcgccga gctgggtcaaa gaggtcgcca agaagaccga	60
cgatgtcgcg ggtgacggca ccaccaccgc caccgtcctg gccagggcgc tgggtgcgca	120
gggtctgcgc aacgttgccg ccggcgccaa cccgatggct ctgaagcgcg gtatcgagaa	180
ggccgtcgag aaggtcaccg agaccctgct gaagtcggcc aaggaggctg agaccaagga	240
gcagatcgct tcgaccgccg cgatctcggc cggcgacacc cagatcggcg agctgatcgc	300
cgaggccatg gacaagg tgc gcaacgagg tgatcatcacc gtcgaggaga gcaacacctt	360
cggcctgcag ctggagctca ccgagggtat gcgcttcgac aagggttaca tctcgggtta	420
cttcgtgacc gacgccgagc gtcaggaagc cgtcctcgag gatccgtaca tcctgctggt	480
gtcgggcaag gtctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc	540
gggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggccctgt cgaccctggt	600

Sequence Listing

cgtg 604

<210> 35

<211> 604

<212> DNA

<213> *Mycobacterium peregrinum*

<400> 35

ggaggacccg tacgagaaga tcggcgctga gctggtcaaa gaggtcgcca agaagaccga 60

cgacgtcgcg ggtgacggca ccaccaccgc caccgttctg gcccaggccc tggttcgcg 120

aggtctgcgc aacgtcgctg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa 180

ggctgtcgag aaggtcaccg agaccctcct gaagtccgcc aaggaggtgg agaccaagga 240

gcagatcgct gccaccgccg gtatctccgc cggagaccag tccatcggcg acctgatcgc 300

cgaggccatg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggaga gcaacacctt 360

cgggctgcag ctggagctca ccgagggcat gcgcttcgac aagggtaca tctcgggcta 420

cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gatccctaca tcctgctggt 480

cagctcgaag atctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc 540

cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctgt cgaccctggt 600

ggtc 604

<210> 36

<211> 604

<212> DNA

<213> *Mycobacterium scrofulaceum*

<400> 36

ggaggacccg tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga 60

Sequence Listing

cgacgtcgcc ggtgacggca cgacgacggc cacggtgctg gcccaggcgc tggtaagga	120
gggcctgcgc aacgtcgcg cgggcgccaa cccgctgagc ctcaagcgcg gcatcgagaa	180
ggcggtcgag aaggtcaccg agaccctgct caagtcggcc aaggaggctg agaccaagga	240
ccagatcgcc gccaccgcg cgatttcggc gggcgaccag tcgatcggcg acctgatcgc	300
cgaggcgatg gacaaggctg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgaggcat gcggttcgac aagggtctaca tctcgggcta	420
cttcgtcacc gacgccgagc ggcaggaagc ggtcctggag gacccttaca tctgctggt	480
cagctcgaag gtgtcgacgg tcaaggacct gctgccgctg ttggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgcttt ccaccctggt	600
cgtc	604

<210> 37
 <211> 604
 <212> DNA
 <213> *Mycobacterium senegalense*

<400> 37	
ggaggaccgc tacgagaaga tcggcgctga gctggtcaag gaagtcgcca agaagactga	60
cgacgtcgcg ggtgacggca ccaccaccgc caccgttctg gcccaggccc tggttcgtga	120
aggtctcgct aacgtcgctg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggccgctcgag aaggtcaccg agacgctgct caagagcgcc aaggaggtag agaccaagga	240
gcagatcgct gccaccgccc cgatctcggc gggcgacacc cagatcgcca agctgatcgc	300
cgaggccatg gacaaggctg gcaacgaggg tgcatcacc gttgaggagt ccaacacctt	360
cgggctgcag ctcgagctca ccgagggtat gcggttcgac aagggtctaca tctcgggtta	420

Sequence Listing

cttcgtgacc gacgccgagc gtcaggaagc ggtcctcgag gatccctgca tcctgctcgt	480
gtcgtccaag gtgtcgaccg tcaaggatct gctcccgttg ctggagaagg tcattcaggc	540
cggcaagccg gtgctgatca tcgccgagga cgtcgagggc gaggccctgt cgaccctggt	600
ggtc	604
<div style="display: flex; justify-content: space-between;"> <210> 38 <211> 604 </div> <div style="display: flex; justify-content: space-between;"> <212> DNA <213> Mycobacterium shimoidei </div>	
<div style="display: flex; justify-content: space-between;"> <400> 38 </div>	
ggaggacccg tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga	60
cgacgtcgcc ggtgacggca ccaccaccgc caccgtgctg gcccaggcgc tggtcacga	120
ggggctgcgc aacgtcgcg ccggtgccaa cccgctcagc ctgaaacgcg gtatcgagaa	180
ggccgttgag aaggtcaccg agaccttgct caaggcgcc aaggaagtcg agaccaagga	240
gcagatcgcg gccacggcgg ccatctccgc cggtgaccag tcgatcggcg acctgatcgc	300
cgaggcgatg gacaaggctg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgagggat gcggttcgac aagggttaca ttccgggtta	420
cttcgtcacc gacgccgagc gtcaggaggc tgtgctcgag gagccctaca tcctgctggt	480
cagctccaag gtgtcgaccg tcaaggacct gctgccgctg ctggagaagg tcatgcaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggctttga gcaccctggt	600
cgtc	604
<div style="display: flex; justify-content: space-between;"> <210> 39 </div>	

Sequence Listing

<211> 604
 <212> DNA
 <213> *Mycobacterium simiae*

<400> 39
 ggaggacccc tacgagaaga tcggcgctga gctggtcaag gaagtcgcca agaagaccga 60
 cgacgtcgcc ggtgacggca ccacgacggc caccgtgctc gctcaggcgc tcgtcaagga 120
 gggcctgcgc aacgtggcgg ccggcgccaa cccgctgggc ctcaagcgcg gcatcgagaa 180
 ggccgtcgaa aaggtcaccg agacgtgct gaagtcggcc aaggatgtcg agaccaagga 240
 ccagatcgct gccaccgccg cgatttccgc gggcgaccag tcgatcggcg acctgatcgc 300
 cgaggcgatg gacaaggctg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt 360
 cgggctgcag ctcgagctca ccgagggat ggcgttcgac aagggtctaca tctcgggcta 420
 cttcgtcacc gacgccgagc gtcaggaagc cgtcctggag gaccggttca tcctgctggt 480
 cagctccaag gtgtcgacgg tcaaggacct gctgccgctg ctggagaagg tcatccaggc 540
 cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctga gcaccctggt 600
 cgtc 604

<210> 40
 <211> 604
 <212> DNA
 <213> *Mycobacterium smegmatis*

<400> 40
 cgaggacccc tacgagaaga tcggtgctga gctcgtcaaa gaggtcgcca agaagaccga 60
 cgatgtcgct ggcgacggca ccaccaccgc caccgtcctg gctcaggccc tggttcgca 120
 aggcctgcgc aacgtcgctg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa 180

Sequence Listing

ggccgctcgag aaggtcaccg agaccctgct gaagtcgcc aaggaggtgg agaccaagga	240
gcagatcgct gccaccgccg gtatctccgc cggtgaccag tccatcggcg acctgatcgc	300
cgaggccatg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgagggat ggccttcgac aagggttaca tctcgggtta	420
cttcgtgacc gacgccgagc gtcaggaagc ggtcctcgag gatccctaca tcctgctggt	480
cagctcgaag gtctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctgt cgaccctggt	600
ggtc	604

<210> 41
 <211> 604
 <212> DNA
 <213> Mycobacterium szulgai

<400> 41	
ggaggaccg tacgagaaga tcggcgccga gctggtcaag gaagttgcca agaagaccga	60
cgacgtcgcc ggtgacggca cgacgacggc caccgtgttg gccaggcgcg tggtaagga	120
ggcctgcgc aacgtagcgg ccggcgccaa cccgctgggt ctcaagcgcg gcatcgagaa	180
ggcgtcgag aagatcaccg agaccctgct caagtcggct aaggacgtcg agaccaagga	240
gcagatcgcg gccaccgccg ccatctccgc gggcgaccag tcgatcggcg acttgatcgc	300
cgaggcgatg gacaaggtcg gcaatgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
cggcctgcag ctcgagctca ccgagggcat gcggttcgac aagggttaca tctcgggcta	420
cttcgtcacc gacgccgagc gtcaggaggc cgtcctcgag gacccttaca tcctgttggt	480
cgctccaag gtgtcgacgg tcaaggacct gttgccgctg ctggagaagg tcatccaggg	540

Sequence Listing

cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggctttga gcaccctggt 600

cgtc 604

<210> 42

<211> 604

<212> DNA

<213> *Mycobacterium terrae*

<400> 42

ggaggacccc tacgagaaga tcggcgccga gctggtcaaa gaggtcgcca agaagaccga 60

cgatgtcgcc ggtgacggca ccaccacggc caccgtgctg gcacaggcgc tggtaagga 120

aggcctgcgc aacgtggccg ccggcgccaa cccgctggcc ctgaagcgcg gcatcgagaa 180

ggccgtcgag aaggtctccg agaccctgct gaaggacgcc aaggaggtcg agaccaagga 240

gcagatcgcg gctaccgccc ggatctccgc gggcgaccag tccatcggtg acctgatcgc 300

cgaggcgatg gacaaggctg gcaacgaggg tgtcatcacc gtcgaggagt ccaacacctt 360

cggcctgcag ctggagctca ccgaggggtat gcgcttcgac aagggtctaca tctcgggtta 420

cttcgtcacc gacgccgacc gtcaggaagc ggttctcgag gaccctaca tcctgctggt 480

cagctccaag atctcgacgg tcaaggacct gctcccactg ctggagaagg tcattcaggg 540

cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggccctgt ccaccctggt 600

ggtc 604

<210> 43

<211> 604

<212> DNA

<213> *Mycobacterium thermoresistibile*

Sequence Listing

<400> 43

ggaggacccc tacgagaaga tcggcgctga gctgggtcaag gaagtcgcca agaagaccga 60

cgacgtcgcc ggcgacggca ccaccaccgc caccgtcctg gctcaggcgc tggatgaagga 120

aggtttgcgc aacgtcgccg ccggggccaa cccgctcgct ctgaagcgcg gcatcggagc 180

cgctgtcgag aaggtcaccg agaccctgct caagtcggcc aaggaggctg agaccaagga 240

gcagatcgcc aacaccgccg cgatctcgcc cggcgaccag cagaccggtg agctgatcgc 300

cgaggcgatg gacaaggctg gcaacgaggg tgtcatcacc gtcgaggagt cgcagacctt 360

cggtctgcag ctcgagctca ccgagggtat gcgcttcgac aagggtctaca tctcggggta 420

cttcgtgacc gacgcggagc ggcaggaagc cgttctggag gatccctaca tcctgctggt 480

cagctcgaag gtctcgactg tcaaggatct gctgccgtg ctggagaagg tcatccagtc 540

cggcaggccg ctgctgatca tcgccgagga cgtcgaaggc gaggcgctgt cgaccctggt 600

cgtc 604

<210> 44

<211> 604

<212> DNA

<213> *Mycobacterium triviale*

<400> 44

ggaggacccg tacgagaaga tcggcgccga gctgggtcaag gaagtcgcca agaagaccga 60

cgatgtcgcc ggtgacggca ccaccacggc caccgtgctc gccagggcgc tggatgcgca 120

gggcctgcgc aacgtcgccg ccggcgccaa cccgatgggc ctgaagcgcg gcatcggaggc 180

ggccaccgag aagatcgccg agaccctgct caaggcgcc aaagagggtg agaccaagga 240

gcagatcgct gccaccgccg ggatctccgc cggggacagc tccatcgggtg agctgatcgc 300

Sequence Listing

cgaggcgatg gacaagggtcg gcaacgaggg tgtcatcacc gtcgaggagg cccagacctt	360
cggcctgcag ctcgagctca ccgaggggtat gcgcttcgac aagggtctaca tctccggcta	420
cttcgtcacc gacgccgagc gtcaggaggc cgtgctggag gaccctaca tcctgctggt	480
gtccggcaag gtgtccaccg tcaaggacct gcttcgctg ctggagaagg tcatccagtc	540
cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt cgaccctggt	600
ggtc	604

<210> 45
 <211> 604
 <212> DNA
 <213> Mycobacterium tuberculosis

<400> 45 ggaggatccg tacgagaaga tcggcgccga gctggtcaaa gaggtagcca agaagaccga	60
tgacgtcgcc ggtgacggca ccacgacggc caccgtgctg gccaggcgt tggttcgga	120
gggcctgcgc aacgtcgcg ccggcgccaa cccgctcggt ctcaaacgcg gcatcgaaaa	180
ggccgtggag aagggtaccg agaccctgct caagggcgcc aaggaggtcg agaccaagga	240
gcagattgcg gccaccgcag cgatttcggc gggtgaccag tccatcggtg acctgatcg	300
cgaggcgatg gacaagggtg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt	360
tgggctgcag ctcgagctca ccgaggggtat gcggttcgac aagggtctaca tctcgggta	420
cttcgtgacc gaccggagc gtcaggaggc ggtcctggag gaccctaca tcctgctggt	480
cagctccaag gtgtccactg tcaaggatct gctgccgctg ctcgagaagg tcatcgagc	540
cggaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt	600
cgtc	604

Sequence Listing

<210> 46
 <211> 604
 <212> DNA
 <213> *Mycobacterium ulcerans*

<400> 46
 ggaggacccg tacgagaaga ttggcgctga gctgggtcaag gaagttgcc aagaagaccga 60
 cgacgtggcc ggtgacggca cgacgacggc caccgtgctg gccagggcg tggtcaagga 120
 aggcctgcgc aacgttgccg ccggtgccaa cccgctcggc ctgaagcgcg gcatcgagaa 180
 ggcagtcgag aaggtcaccg agaccctgct caaatcggcc aaagaggtcg agaccaagga 240
 gcagatcgcg gcgaccgag ccatctccgc cggcgaccag tcgatcggcg acctgatcgc 300
 cgaggcgatg gacaaggtgg gcaacgaggg cgtcatcacc gtcgaggagt ccaacacctt 360
 cggcctgcag ctcgagctca ccgaggggat gcggttcgac aagggtctaca tctcgggcta 420
 cttcgtcacc gacgcccagc gtcaggaagc ggtcctggag gaccctaca tcctgctggt 480
 cagctccaag gtgtccaccg tcaaggacct gctgccgctg ctggagaagg tcattcaggg 540
 cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggcgctgt ccaccctggt 600
 cgtc 604

<210> 47
 <211> 604
 <212> DNA
 <213> *Mycobacterium vaccae*

<400> 47
 ggaggacccg tacgagaaga tcggcgctga gctgggtcaaa gaggtcgcca aagaagaccga 60
 cgacgtcgcg ggcgacggta ccaccaaccg caccgtgctc gtcaggtc tggttcgga 120

Sequence Listing

aggcctgcgc aacgtcgcag ccggcgccaa cccgctcggc ctcaagcgtg gcatcgagaa	180
ggctgtcgag gctgtcacc agtcgtgct gaagtcggcc aaggaggtcg agaccaagga	240
gcagatttct gccaccgcg cgatctccgc cggcgacacc cagatcggcg agctcatcgc	300
cgaggccatg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggagt cgaacacctt	360
cggcctgcag ctcgagctca ccgagggat gcgcttcgac aagggtaca tctcgggtta	420
cttcgtgacc gacgccgagc gccaggaagc cgtcctggag gatccctaca tcctgctggt	480
cagctccaag gtgtcgaccg tcaaggatct gtcctcgctg ctggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggccctgt ccacgctggt	600
ggtc	604

<210> 48
 <211> 604
 <212> DNA
 <213> Mycobacterium wolinskyi

<400> 48 ggaggacccg tacgagaaga tcggcgctga gctgggtcaaa gaggtcgcca agaagaccga	60
cgacgtcgcc ggcgacggca ccaccaccgc caccgttttg gccaggctc tggttcgca	120
aggtctgcgc aacgtcgcgg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggccgtcgag aaggtcaccg agacgtgct gaagagcgcc aaggaggtgg agaccaagga	240
gcagatcgt gccaccgcc gtatctccgc cggtgaccag tccatcggcg acctgatcgc	300
cgaggccatg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggaga gcaacacctt	360
cggcctgcag ctggagctca ccgagggat gcgcttcgac aagggtaca tctcgggtta	420

Sequence Listing

cttcgtgacc gacgccgagc gtcaggaagc cgtcctcgag gatccctaca tcctgctggt	480
cagctcgaag gcctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggccctgt cgaccctggt	600
ggtc	604

<210> 49
 <211> 604
 <212> DNA
 <213> *Mycobacterium parafortuitum*

<400> 49 ggaggacccg tacgagaaga tcggcgctga gctgggtcaaa gaggtcgcca agaagaccga	60
cgacgtcgcg ggcgacggca ccaccaccgc caccgtgctc gctcaggccc tggttcgcg	120
aggtctgcmc aacgtcgcmg ccggcgccaa cccgctcgmc ctcaagcmgm gcatcgagaa	180
ggctgtcmg gctgtcacc agggctctgct gaagtcggcc aaggaggtcm agaccaagga	240
gcagatcmgt gccaccgcmg cmgtctcmg cggcgacacc cmgtcmgm agctcatcmg	300
cmggcccatg gacaaggtcm gcaacgaggg tgtcatcacc gtcgaggagt cgaacacctt	360
cmgcctcmg ctggagctca ccgaaggcat gcgcttcgac aagggtctaca tctcmgggta	420
cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gatccctaca ttctgctggt	480
cagctccaag atctcmagcm tcaaggacct gctgccgctg ctggagaagg tcatccagtc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctgt cgaccctggt	600
ggtc	604

<210> 50
 <211> 604

Sequence Listing

<212> DNA

<213> *Mycobacterium farcinogenes*

<400> 50

ggaggacccg tacgagaaga tcggcgctga gtcgtcaaa gaggtcgcca agaagaccga	60
cgacgtcgcg ggcgacggca ccaccaccgc caccgttctg gcccaggccc tggttcgcg	120
aggtctgcgc aacgtcgctg ccggcgccaa cccgctcggc ctgaagcgcg gcatcgagaa	180
ggccgtcgag aaggtcaccg agacgtgct caagagcgcc aaggaggtgg agaccaagga	240
gcagatcgct gccaccgccg gtatctccgc cggtgaccag tccatcggtg acctgatcgc	300
cgaggccatg gacaaggtcg gcaacgaggg tgtcatcacc gtcgaggaga gcaacacctt	360
cggcctgcag ctggagctca ccgaggggat gcgcttcgac aagggtctaca tctcgggtta	420
cttcgtgacc gacgccgagc gtcaggaagc cgtcctggag gatccctaca tctgctggt	480
cagctccaag gtctcgaccg tcaaggatct gctgccgctg ctggagaagg tcatccagtc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaagccctgt ccaccctggt	600
ggtc	604

<210> 51

<211> 604

<212> DNA

<213> *Tsukamurella paurometabola*

<400> 51

cgaggatccc tacgagaaga tcggcgccga gtcgtcaag gaggtcgcca agaagaccga	60
cgacgtcgcg ggcgacggca ccaccaccgc caccgttctg gcccaggcgc tcgtcgcgca	120
gggtctgcgc aacgtggctg cgggtgcgaa cccgctgggc ctcaagcggg gcatcgagaa	180
ggccgtcgag gccgtgaccg agcacctgct caaggaggcc aaggaggtcg agaccaagga	240

Sequence Listing

gcagatcgct gctaccgcgg gcatctcggc cggcgacccc gccatcggtg agtcatcgc	300
cgaggccatg gacaaggctg gcaaggaagg cgtcatcacc gtcgaggaga gcaacacctt	360
cggctctccag ctggagctca ccgagggcat gcgcttcgac aagggttca tctcggcta	420
cttcgccacc gacgccgagc gtcaggaggc cgtgctcgag gacgcctaca tcctgctcgt	480
gtcgagcaag atctcgaccg tgaaggacct gctgccgctg ctggagaagg tcatccagtc	540
gggcaagccg ctcgcatca tcgccgagga cgtcgagggc gaggccctgt cgacgctcat	600
cgtc	604

<210> 52
 <211> 604
 <212> DNA
 <213> Tsukamurella tyrosinosolvens

<400> 52	
cgaggatccc tacgagaaga tcggcgccga gctcgtcaag gaggtcgcca agaagaccga	60
cgacgtcgcg ggcgacggca ccaccaccgc caccgttctg gccaggcgc tcgtgcgca	120
gggcctgcgc aacgtggccg cggcgcgaa cccgctgggc ctcaagcggg gcatcgagaa	180
ggccgtcgag gccgtctccg agcacctgct gaaggccgcc aaggaggtcg agaccaagga	240
gcagatcgct gctaccgcgg gcatctcggc cggcgacccc gccatcggtg agtcatcgc	300
cgaggccatg gacaaggctg gcaaggaagg cgtcatcacc gtcgaggaga gcaacacctt	360
cggcctccag ctggagctca ccgagggcat gcgcttcgac aagggttca tctcgggcta	420
cttcgccacc gacgccgagc gtcaggaggc cgtgctcgag gacgcctacg tgctgctcgt	480
cgccggcaag atctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc	540

Sequence Listing

gggcaagccg ctcgcatca tcgccgagga cgtcgagggc gaggccctgt cgacgctcat 600

cgtc 604

<210> 53

<211> 604

<212> DNA

<213> *Tsukamurella pulmonis*

<400> 53

cgaggatccc tacgagaaga tcggcgccga gctcgtcaag gaggtcgcca agaagaccga 60

cgacgtcgcg ggcgacggca ccaccaccgc caccgttctg gccaggcgc tcgtgcgcga 120

gggtctgcgg aacgtggccg cgggcgcgaa cccgctgggc ctcaagcggg gcatcgagaa 180

ggcggtcgac gccgtcaccg agcacctgct gaaggccgcc aaggaggtcg agaccaagga 240

gcagatcgct gctaccgcgg gcatctcggc cggcgacccc gccatcgggtg agctcatcgc 300

cgaggccatg gacaaggctg gcgaggaagg cgtcatcacc gtcgaggaga gcaacacctt 360

cggctctccag ctggagctga ccgagggcat gcgcttcgac aagggttca tctcgggcta 420

cttcgccacc gacgcggagc gccaggaggc cgtcctcgag gacgcctacg tgctgctcgt 480

ctcgggcaag atctcgaccg tcaaggacct gctgccgctg ctggagaagg tcatccagtc 540

gggcaagccg ctcgcatca tcgccgagga cgtcgagggc gaggccctgt cgacgctcat 600

cgtc 604

<210> 54

<211> 604

<212> DNA

<213> *Nocardia carnea*

<400> 54

Sequence Listing

cgaggatccc tacgagaaga tcggcgccga gctggtcaag gaagtcgcca agaagaccga	60
cgacgtcgcg ggcgacggca ccaccaccgc caccgtgctc gcccagggcg tggtgcgcga	120
gggtctgcgc aacgtggccg cgggcgcgaa cccgctgggc ctcaagcgca gcatcgagaa	180
ggccgtcgag gccgtgaccg ccaagctgct cgacaccgcc aaggaggtcg agaccaagga	240
gcagatcgcc gccaccgcgg gcatctccgc gggcgacgcg tccatcggtg agctgatcgc	300
cgaggccatg gacaaggctg gcaaggaagg cgtcatcacc gtcgaggaga gcaacacctt	360
cggcctccag ctggagctga ccgagggcat gcgcttcgac aagggctaca tctccggcta	420
cttcgtgacc gatcccgagc gtcaggaagc ggtcctcgag gatccctaca tctgctcgt	480
cggctcgaag gtctccaccg tcaaggacct gctgccgctg ctggagaagg tcatccaggc	540
cggcaagccg ctgctgatca tcgccgagga cgtcgagggc gaggccctgt cgaccctggt	600
cgtg	604

<210> 55
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> HSPF3

<400> 55	
atcgccaagg agatcgagct	20

<210> 56
 <211> 20
 <212> DNA
 <213> Artificial Sequence

Sequence Listing

<220>

<223> HSPR3

<400> 56

aaggtgccgc ggatcttggt

20

INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR03/00131

A. CLASSIFICATION OF SUBJECT MATTER

IPC7 C12Q 1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 : C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

REGISTRY/STN, PubMed

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Ena P. et al. "Rapid identification of cutaneous infections by nontubercular mycobacteria by polymerase chain reaction-restriction analysis length polymorphism of the hsp65 gene." In: Int. J. Dermatol., 2001 Aug., 40(8): pages 495-499, see entire document.	1-16
A	Yakrus M.A. et al. "Comparison of methods for identification of Mycobacterium abscessus and M. chelonae isolates." In: J. Clin. Microbiol., 2001 Nov., 39(11): pages 4103-4110, see entire document.	1-16
A	Goh K.S. et al. "Rapid differentiation of Mycobacterium canettii from other Mycobacterium tuberculosis complex organisms by PCR-restriction analysis of the hsp65 gene." In: J. Clin. Microbiol., 2001 Oct., 39(10): pages 3705-3708, see entire document.	1-16
A	Brunello F. et al. "Identification of 54 mycobacterial species by PCR-restriction fragment length polymorphism analysis of the hsp65 gene." In: J. Clin. Microbiol., 2001 Aug., 39(8): pages 2799-2806, see entire document.	1-16
A	Leclerc M.C. et al. "Molecular characterization of environmental Mycobacterium strains by PCR-restriction fragment length polymorphism of hsp65 and by sequencing of hsp65, and of 16S and ITS1 rDNA." In: Res. Microbiol., 2000 Oct., 151(8): pages 629-638, see entire document.	1-16

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 MAY 2003 (23.05.2003)

Date of mailing of the international search report

24 MAY 2003 (24.05.2003)

Name and mailing address of the ISA/KR



Korean Intellectual Property Office
920 Dunsan-dong, Seo-gu, Daejeon 302-701,
Republic of Korea

Facsimile No. 82-42-472-7140

Authorized officer

HAN, Hyung Mee

Telephone No. 82-42-481-5601

